

Phylogenetic analysis and antibiotic resistance of *Shigella sonnei* isolates

Babak Pakbin^{1,2,3}, Abdollah Didban², Wolfram M. Brück^{1,*}, Mehdi Alizadeh²

¹Institute for Life Technologies, University of Applied Sciences Western Switzerland Valais-Wallis, 1950 Sion 2, Switzerland

²Children Growth Research Center, Research Institute for Prevention of Non-Communicable Diseases, Qazvin University of Medical Sciences, Bahonar Blvd., Qazvin 34185–754, Iran

³Medical Microbiology Research Center, Qazvin University of Medical Sciences, Qazvin 34197–59811, Iran

*Corresponding author: Institute for Life Technologies, University of Applied Sciences Western Switzerland Valais-Wallis, 1950 Sion 2, Switzerland.

Tel: (+41) 586068664; E-mail: wolfram.bruck@hevs.ch

One sentence summary: We determined the phylogeny and antibiotic resistance profiles of *S. sonnei* strains and found most of the strains clustered within the lineage III and resistance against trimethoprim-sulfamethoxazole and tetracycline antibiotics.

Editor: Rustam Aminov

Abstract

Shigellosis is one of the most important gastric infections caused by different species of *Shigella*, and has been regarded as a serious threat to public health. Lineage/sublineage profile of *Shigella sonnei* is strongly associated with the antibiotic resistance and population structure of this pathogen. In this study, we determined the phylogeny and antibiotic resistance profiles of *S. sonnei* strains, isolated from 1246 stool and 580 food samples, using multiplex PCR–HRMA genotyping and Kirby–Bauer disk diffusion methods, respectively. A total of 64 *S. sonnei* strains were isolated (13 food and 51 clinical isolates). Multiplex PCR–HMR assay was able to differentiate the lineages II and III, and sublineages IIIb and IIIc strains successfully considering the definite melting curves and temperatures. Lineage I and sublineage IIIa strain were not isolated in this study. We also demonstrated that most of the *S. sonnei* strains isolated from both food and clinical samples clustered within the lineage III and sublineage IIIc. Resistance against trimethoprim-sulfamethoxazole, tetracycline, chloramphenicol, and streptomycin antibiotics were the most prevalent phenotypes among the *S. sonnei* lineage III and sublineage IIIc strains.

Keywords: *Shigella sonnei*, PCR–HMRA, phylogenetic group, antibiotic resistance

Introduction

Shigella sonnei is a Gram-negative, rod-shaped, and facultative intracellular pathogen belonging to *Shigella* and *Enterobacteriaceae* family (Pakbin et al. 2021c; Torraca et al. 2020). *Shigella sonnei* has been regarded as the main cause of shigellosis in developed countries in North America and Europe. Also, *S. sonnei* has frequently been isolated from shigellosis in those that are economically damaged or developing in the high-income countries (Shad and Shad 2021). However, it is currently expanding in some middle-income countries across the Middle East, Latin America, and Asia (Baker and The 2018). This bacterial pathogen causes self-limiting and acute gastrointestinal diseases characterized by fever, watery or bloody diarrhoea, and abdominal pain symptoms in adults and children. The infection caused by this pathogen can be strongly life-threatening for infants and children under 5 years old (Rogawski McQuade et al. 2020). It is estimated that this pathogen causes annually more than 188 million cases and 160 000 deaths, mainly among young children (Hosangadi et al. 2019). *Shigella sonnei* has been considered a foodborne pathogen, and commonly spreads via faecal–oral transmission as it can survive gastric acidity and infects the colonic mucosa (Belotserkovsky and Sansonetti 2018). The main treatment for shigellosis is rehydration therapy and antibiotics such as cephalosporins, fluoroquinolones, and azithromycin. However, resistance against different antibiotics is becoming more common among the *S. sonnei* isolated from

food and clinical samples. It is widely reported that some specific genotypes of *S. sonnei* are associated with high rates of multidrug resistance and more frequently contribute to foodborne outbreaks (Pakbin et al. 2020a, Pakbin et al. 2021a; Ranjbar and Farahani 2019).

The genotypic profile of *S. sonnei* isolates is strongly associated with the pathogenicity, resistance against different antibiotics, regional epidemiology, and population structure of the pathogen (Pakbin et al. 2021b, c; The et al. 2021). Consequently, genotype characterization of *S. sonnei* isolates helps to better investigate the outbreaks in respect of the presence of this pathogen and choose the most efficient strategies to prevent the spread of shigellosis due to the *S. sonnei* isolates. *Shigella sonnei* is monoclonal, and there is only one serogroup (D) consisting of a single serotype (Ragupathi et al. 2018). Whole-genome sequencing (WGS) has effectively been used for subtyping the *S. sonnei* isolates. According to the WGS studies, *S. sonnei* is categorized into five significant lineages and several sublineage phylogenetic groups associated with some phenotypic features, including resistance to different antibiotics, the transmission routes, and regional epidemiology (Dallman et al. 2016). For instance, *S. sonnei* isolated from the specimen in Australia, Europe, North America, and Asia have been dominated by the Lineage 3 strains for more than 29 years (Holt et al. 2012). These strains demonstrated resistance against tetracycline, trimethoprim-sulfamethoxazole, streptomycin, ampicillin,

Received: February 9, 2022. Revised: March 28, 2022. Accepted: April 26, 2022

© The Author(s) 2022. Published by Oxford University Press on behalf of FEMS. All rights reserved. For permissions, please e-mail: journals.permissions@oup.com

chloramphenicol, ciprofloxacin, and fluoroquinolones in the presence of antibiotic resistance genes and through the acquisition of point mutations (Hawkey et al. 2021). A single nucleotide polymorphisms (SNPs)-typing scheme based on the high-resolution melting analysis (HRMA) method was developed and described for the first time by Sangal et al. (2013) to identify all lineages and sublineages of *S. sonnei* strains. They detected SNPs in small PCR amplicons (SNPs within *deoA*, *kduD*, and *emrA* genes for typing three main lineages I, II, and III, and SNPs within *menF* and *fdx* genes for typing sublineages including IIIa, IIIb, and IIIc) based on the high-resolution melting patterns. They reported this method is more rapid and cost-effective than WGS for genotyping *S. sonnei* strains (Sangal et al. 2013). Mazi et al. (2015) developed a multiplex PCR–HRMA method to identify and differentiate the lineages I, II, III and sublineages IIIa, IIIb, and IIIc of the *S. sonnei* strains using the Qiagen Rotor-Gene 6000 instrument and EvaGreen dye (Mazi et al. 2015). Since the characterization of genotype, lineage, sublineages, and antibiotic resistance of *S. sonnei* strains represents an opportunity to understand the ecology, epidemiology, and antibiotic resistance of this pathogen to choose the best preventive strategies (Hawkey et al. 2021), this study aimed to determine the lineages, sublineages, and antibiotic resistance profile of *S. sonnei* strains isolated from clinical and food samples by using the PCR–HRMA method as a rapid, precise, and cost-effective assays.

Materials and methods

Sample collection

Stool samples ($N = 1246$) from children and adults with acute diarrhoea referred to the emergency service and central lab in different hospitals (Qazvin children hospital, Qazvin, Iran and Milad hospital, Tehran, Iran), and the food samples ($N = 580$), including raw milk ($n = 110$), ground meat ($n = 210$), and raw vegetable ($n = 260$) samples from local markets located in various areas of municipalities of Tehran and Qazvin, Iran were collected during Jun 2015 to August 2018. All samples were immediately transported in cool boxes ($4^{\circ}\text{C} \pm 0.5$) to the central microbiology research laboratory for further analysis. *Shigella sonnei* PTCC 1777 was purchased from Pasteur Institute (Pasteur In., Tehran, Iran) and used as the positive control and reference strain in this study.

Isolation and identification of *S. sonnei*

Shigella sonnei was isolated and identified in stool and food samples according to the methods described by Phiri et al. (2021) and Ahmed and Shimamoto (2014), respectively. Suspected colonies (red) on XLD agar were selected and subjected to the biochemical tests, including IMViC, triple sugar iron (TSI), urease production, and lysine decarboxylase tests. Slide agglutination method using *Shigella* genus and species polyvalent antisera commercial kits (Difco Co., MI, USA) was used according to the manufacturers' instructions to confirm and determine the genus and species of the presumptive colonies, respectively. All strains were subjected to DNA extraction.

Antibiotic susceptibility testing

Antibiotic resistance of *S. sonnei* isolates was performed using Kirby–Bauer disk diffusion method based on the interpretive criteria and standards previously developed and described by Clinical and Laboratory Standards Institute (Pakbin et al. 2020b). A total of 14 commercial antibiotic disks (Oxoid, UK) used in this study included kanamycin (KAN), 30 μg ; amoxicillin-clavulanic acid (AMC), 20/10 μg ; streptomycin, 10 μg ; ampicillin (AMP), 10 μg ;

Table 1. The primers used in this study for PCR–HRM genotyping of *S. sonnei* isolates.

Target gene	Primer	Sequence (5'→3')
<i>kduD</i>	kduDF	CGACGGCGAAACACTTTTATC
	kduDR	CGCGTATAAGAAGGCACACG
<i>emrA</i>	emrAF	TGCCACCGAAGTACGTAACG
	emrAR	CATCCACCCACATATTGGTG
<i>deoA</i>	deoAF	GGAGATGCTTATCTCCGGCAAA
	deoAR	AGTCGGTTGGGCCTTTT
<i>menF</i>	menFF	TATTTCTCGCGCTGGTTTTTA
	menFR	GCTTTTCTGGCTCTTACC
<i>fdx</i>	fdxF	CAAAGCCTGGGACTGGA
	fdxR	CATGGTTGATAGTGTAAACG

amoxicillin (AMX), 25 μg ; amikacin (AMK), 30 μg ; azithromycin (AZM), 15 μg ; gentamicin (GEN), 10 μg ; colistin (CST) 10 μg ; chloramphenicol (CHL), 30 μg ; trimethoprim-sulfamethoxazole (SXT), 1.25/23.75 μg ; tetracycline (TET), 30 μg ; nalidixic acid (NAL), 30 μg ; levofloxacin (LVX), 5 μg , and nitrofurantoin (NIT), 300 μg . The results of antibiotic resistance phenotypes were recorded and interpreted according to CLSI guidelines (Pakbin et al. 2020b). *Klebsiella pneumoniae* ATCC 700603, *Staphylococcus aureus* ATCC 25923, and *Escherichia coli* ATCC 25922 were used as the reference strains in this study.

DNA extraction

According to the manufacturers' instructions, the genomic of the bacterial strains was extracted using the Sinaclon Gram-negative bacterial DNA extraction commercial kit (Sinaclon Co., Tehran, Iran). The quantity and quality of the extracted DNA were assessed using the NanoDrop-1000 spectrophotometer (ThermoFisher, MD, USA). Final concentrations of the extracted genomes were adjusted to 50 $\text{ng } \mu\text{l}^{-1}$. All DNA templates were kept at -20°C until genotyping analysis.

Genotyping by PCR–HRMA

Lineages and sublineages of *S. sonnei* strains isolated from food and clinical samples in this study were determined and differentiated using a PCR–HRMA method previously described by Mazi et al. (2015). Primer sequences used at the present study for PCR–HRMA genotyping of the isolates are summarized in Table 1. PCR–HRMA was performed using a Rotor-Gene Q 6000 real-time PCR instrument (Corbett, Australia) in this study. Genotyping of the strains was performed in two separate runs. In the first run, each reaction contained 20 μl including 4 μl of 5X EvaGreen hot-start PCR–HRM master mix (Solis BioDyne Co, Estonia), 2 μl of DNA template (10 $\text{ng } \mu\text{l}^{-1}$), 0.5 μl of each primer of *emrA* (100 $\mu\text{M } \mu\text{l}^{-1}$), 1 μl of each primer of *deoA* (100 $\mu\text{M } \mu\text{l}^{-1}$), and 2.5 μl of each primer of *kduD* (100 $\mu\text{M } \mu\text{l}^{-1}$) and nuclease-free water up to the final reaction volume. In the second set, PCR–HRM was performed in a 20- μl final reaction volume containing 4 μl of 5X EvaGreen hot-start PCR–HRM master mix (Solis BioDyne Co, Estonia), 2 μl of DNA template (10 $\text{ng } \mu\text{l}^{-1}$), 2 μl of each primer of *fdx* (10 $\mu\text{M } \mu\text{l}^{-1}$), 1 μl of each primer of *menF* (10 $\mu\text{M } \mu\text{l}^{-1}$), and distilled deionized water up to the final reaction volume. The PCR program for both runs was carried out as follow: initial denaturation at 95°C for 12 min, followed by 35 cycles containing 20 s at 95°C , 20 s at 60°C , and 22 s at 72°C . HRM was conducted between 75 to 95°C , with a data acquisition every 0.1°C . Rotor-Gene 6000 software version 2.02 (Corbett, Australia) was used to analyze the HRM data.

Table 2. T_m profile for each *S. sonnei* lineage detected in this study.

Lineage	T_m ($^{\circ}\text{C}$) ($\pm 0.02^{\circ}\text{C}$)				
	<i>kduD</i>	<i>deoA</i>	<i>emrA</i>	<i>fdx</i>	<i>menF</i>
Main lineages					
II	85.8	88.6	91.1		
III	85.8	88.9	90.9		
Sublineages					
IIIb				85.9	89.4
IIIc				86.3	88.6

Statistical analysis

Chi-square and Fisher's exact tests were used to evaluate the significant differences ($P < .05$) between the groups using the SPSS version 21.0.1 (IBM Corp., NY, USA) software.

Results

Lineages and sublineages of *S. sonnei* strains by using PCR–HRMA

A total of 64 *S. sonnei* strains were isolated and confirmed from 1246 stool and 580 food samples. A total of 20.3% ($n = 13$) and 79.6% ($n = 51$) of all strains were isolated from food and clinical samples, respectively. In order to genotype and define the population structure of *S. sonnei* strains, we used PCR–HRMA method previously developed and optimized by Sangal et al. (2013) and Mazi et al. (2015), respectively. As it is shown in Fig. 1(A), the first HRMA-genotyping set run was clearly and successfully able to separate and differentiate lineages II and III regarding the distinct melting curves. Lineage I was not detected and identified among the strains. The second HRMA-genotyping set run completely distinguished the sublineages IIIb and IIIc with significantly distinct melting curves (Fig. 1B). Sublineage IIIa was not detected also among the *S. sonnei* strains. Melting temperatures of all alleles are summarized in Table 2. Significant differences ($P < .01$) were observed among the melting temperatures of all alleles. Sequencing of all SNPs verified the reliability of the melting temperatures of the amplicons associated with each lineage and sublineage of *S. sonnei* strains as previously described by Mazi et al. (2015). Consequently, lineages and sublineages of *S. sonnei* strains were identified and differentiated correctly in this study.

Lineage and sublineage prevalence of *S. sonnei* strains

Shigella sonnei strains are categorized into three main whole-genome sequence analysis types called 'lineages', which were identified and differentiated by using the PCR–HRMA assay in this study. Phylogenetic analysis of *S. sonnei* strains isolated from food and stool samples were performed to determine its population structure. The prevalence of different *S. sonnei* lineages and sublineages in this study are shown in Fig. 2(A) and (B), respectively. A total of 100% (13 out of 13 isolates) and 96% (49 out of 51 isolates) of *S. sonnei* strains isolated from food and stool samples, respectively clustered within lineage III. Only two strains, which were isolated from stool samples, were divided into lineage II. The strains in lineage III were further divided into the sublineages IIIb and IIIc. A total of 18.3% (9 out of 49) and 81.6% (40 out of 49) *S. sonnei* lineage III strains isolated from stool samples clustered within sublineages IIIb and IIIc, respectively. Most of the *S. sonnei* lineage III strains isolated from food samples (92.3%, 12 out of 13 isolates) were categorized into the sublineage IIIc and only

one food strain (7.6%, 1 out of 13 isolates) was identified the sublineage IIIb. This study did not detect *S. sonnei* strains belonging to lineage I and sublineage IIIa. The majority of *S. sonnei* strains isolated from food and stool samples were significantly ($P < .05$) belonging to the lineage III. Notably, most of the *S. sonnei* lineage III strains isolated from both food and stool samples significantly ($P < .05$) belonged to the sublineage IIIc.

Antibiotic susceptibility profile of *S. sonnei* lineages/sublineages

All 64 *S. sonnei* isolates were evaluated for antibiotic susceptibility against 9 classes and 14 antibiotics. The results of antibiotic susceptibility profiles of different *S. sonnei* lineages and sublineages are presented in Table 3. Trimethoprim-sulfamethoxazole, amoxicillin, tetracycline, ampicillin, amoxicillin-clavulanic acid, streptomycin, and chloramphenicol resistance were dominant phenotypes among the *S. sonnei* strains. However, the lowest antibiotic resistance was observed against kanamycin and nitrofurantoin. *Shigella sonnei* lineage III and sublineage IIIc strains were mostly resistant to trimethoprim-sulfamethoxazole, tetracycline, chloramphenicol, and streptomycin antibiotics.

Discussion

Genotypic characteristics of *S. sonnei* strains are strongly associated with population structure, pathogenicity, and antibiotic resistance properties of this pathogen (Hawkey et al. 2021). Different types of *S. sonnei* have been identified by WGS analysis of this pathogen, categorized into several lineages and sublineages (Baker et al. 2017). Alternative methods such as PCR–HRMA have been designed and developed to differentiate these phylogenetic groups among the *S. sonnei* strains (Mazi et al. 2015). This study used PCR–HRMA assay to identify lineages and sublineages of 64 *S. sonnei* strains isolated from food ($n = 13$) and stool samples ($n = 51$). HRMA method differentiated lineages II and III from each other, considering significantly different melting curves and temperatures. Also, this method determined the sublineages IIIb and IIIc among the strains. The reliability of the melting temperatures of the amplicons was verified by sequencing. Sangal et al. (2013), for the first time, designed and used PCR–HRMA assay for identification of *S. sonnei* phylogenetic groups (lineages/sublineages). They showed that *S. sonnei* phylogeny can be accurately determined with limited SNPs by using this method and different *S. sonnei* lineages/sublineages can be identified successfully as previously defined by the variations of whole-genome (Sangal et al. 2013). Mazi et al. (2015) developed and optimized the multiplex-PCR–HRMA method to differentiate *S. sonnei* lineages and sublineages using Rotor-Gene 6000 HRM real-time PCR instrument (Qiagen) EvaGreen intercalating dye as we used the same dye and instrument in this study. They found that this method is a robust and straightforward genotyping assay for rapid and cost-effective identification of *S. sonnei* lineages and sublineages and might be regarded as an alternative method of WGS (Mazi et al. 2015). The differences between the melting temperatures reported in this and previous studies might be due to the variations in the experimental conditions, such as differences in PCR–HRM master mix and MgCl_2 concentration (Reed et al. 2007; Wittwer 2009).

The results of this study provide novel information about the prevalence of different lineages and sublineages of *S. sonnei* strains isolated from food and stool samples. We found that the prevalence of *S. sonnei* lineage III and sublineage IIIc was significantly higher than that of other lineages and sublineages among the




Figure 1. (A) Multiplex-PCR-HRM curves of lineage II (red curve) and lineage III (blue curve) *S. sonnei* strains; (B) multiplex-PCR-HRM curves of sublineage IIIb (black curve) and sublineage IIIc (Green curve) *S. sonnei* strains.

Table 3. Antibiotic resistance profile of *S. sonnei* lineages/sublineages.

Antibiotic group	Antibiotic agent	n (%)			
		Lineage II (N = 2)	Lineage III (N = 62)	Sublineage IIIb (N = 10)	Sublineage IIIc (N = 52)
β -Lactams	Amoxicillin	2 (100)	37 (59.6)	4 (40.0)	33 (63.4)
	Ampicillin	2 (100)	38 (61.2)	5 (50.0)	33 (63.4)
	Amoxicillin-clavulanic acid	1 (50)	35 (56.4)	7 (70.0)	28 (53.8)
Aminoglycosides	Streptomycin	1 (50)	45 (72.5)	5 (50.0)	40 (76.9)
	Kanamycin	0 (0)	31 (50.0)	2 (20.0)	29 (55.7)
	Amikacin	0 (0)	0 (0)	0 (0)	0 (0)
Quinolones and fluoroquinolones	Nalidixic acid	0 (0)	30 (48.3)	3 (30.0)	27 (51.9)
	Levofloxacin	0 (0)	33 (53.2)	6 (60.0)	30 (57.6)
Macrolides	Azithromycin	0 (0)	32 (51.6)	7 (70.0)	25 (48.0)
Tetracyclines	Tetracycline	1 (50)	49 (79.0)	9 (90.0)	40 (76.9)
Lipopeptides	Colistin	0 (0)	12 (19.3)	2 (20.0)	10 (19.2)
Phenicol	Chloramphenicol	0 (0)	46 (74.1)	8 (80.0)	38 (73.0)
Nitroheterocyclics	Nitrofurantoin	0 (0)	10 (16.1)	3 (30.0)	7 (13.4)
Folate pathway antagonists	Trimethoprim-sulfamethoxazole	1 (50)	54 (87.0)	10 (100)	44 (84.6)





Figure 2. Prevalence rate of different *S. sonnei* strain lineages (A) and sublineages (B) isolated from food and stool samples.

strains isolated from food and clinical samples. Sangal *et al.* (2013) also reported that 58 out of 68 *S. sonnei* isolates (85.2%) belonged to lineage III. They found 41.1% (28 out of 68) and 38.2% (26 out of 68) of the strains belonged to sublineages IIIb and IIIc, which were mostly isolated from Brazil and North Africa, respectively (Sangal *et al.* 2013). Baker *et al.* (2018) investigated the genomic epidemiology of *S. sonnei* isolated from clinical samples in the UK by using the WGS technique, and they also found that most of the *S. sonnei* strains (96.7%; 181 out of 187) belonged to the lineage III (Baker *et al.* 2018). Hawkey *et al.* (2021) studied the global population structure of *S. sonnei* strains and they revealed that most of the *S. sonnei* strains isolated in Asia, Africa, and Latin America are clustered within lineage III, and the sublineages IIIa and IIIb were mostly identified in Central Asia and Latin America (Hawkey *et al.* 2021). This is the first study that investigated the *S. sonnei* lineages/sublineages in food samples, and we found the same results compared to those from the clinical samples. Lineage III was the most common lineage followed by lineage II that researchers have reported among the *S. sonnei* strains worldwide (Baker *et al.* 2017; Hawkey *et al.* 2021; The *et al.* 2021); also, we reported at the present study in Iran among the isolates from food and stool samples.

This study found that resistance against trimethoprim-sulfamethoxazole, tetracycline, chloramphenicol, and streptomycin antibiotics were the dominant phenotypes among the *S. sonnei* strains grouped into the lineage III and sublineage IIIc. As previously described, the strains clustered within lineage III are predominantly resistant to tetracycline, trimethoprim-sulfamethoxazole, and streptomycin because of a chromosomal Tn7-like transposon and the small plasmid spA. A total of 74.1% and 61.2% of *S. sonnei* lineage III isolates in this study were resistant to ampicillin and chloramphenicol antibiotics, respectively. Resistance to ampicillin and chloramphenicol antibiotics mainly was observed among the lineage III strains. Several studies also previously reported resistance against fluoroquinolones and ciprofloxacin among *S. sonnei* lineage III strains due to the acquisition of point mutations within the *gyrA* and *parC* genes (Hawkey *et al.* 2021; Phiri *et al.* 2021; Shad and Shad 2021; The *et al.* 2021). It is extremely important to monitor different *S. sonnei* lineages/sublineages and the antibiotic resistance profile of these strains as the new emerging threat to public health and global concern of food safety (Pakbin *et al.* 2021a,c, Hawkey *et al.* 2021).

Conclusions

This study determined the phylogeny and antibiotic resistance profiles of *S. sonnei* strains isolated from food and stool samples. HMR assay correctly differentiated the lineages II and III, and sublineages IIIb and IIIc strains regarding the distinct melting curves and temperatures. We did not detect any lineage I and sublineage IIIa strain in this study. *Shigella sonnei* strains isolated from both food and clinical samples in this study clustered dominantly within the lineage III and sublineage IIIc. We also found that *S. sonnei* lineage III and sublineage IIIc strains mainly were resistant to trimethoprim-sulfamethoxazole, tetracycline, chloramphenicol, and streptomycin antibiotics.

Acknowledgments

We acknowledge our colleagues at the Medical Microbiology Research Center who contributed to the parts of this work.

Conflicts of interest statement. None declared.

References

- Ahmed AM, Shimamoto T. Isolation and molecular characterization of *Salmonella enterica*, *Escherichia coli* O157: H7 and *Shigella* spp. from meat and dairy products in Egypt. *Int J Food Microbiol* 2014;**168**:57–62.
- Baker K, Campos J, Pichel M *et al.* Whole genome sequencing of *Shigella sonnei* through pulsenet Latin America and Caribbean: advancing global surveillance of foodborne illnesses. *Clin Microbiol Infect* 2017;**23**:845–53.
- Baker KS, Dallman TJ, Field N *et al.* Genomic epidemiology of *Shigella* in the United Kingdom shows transmission of pathogen sublineages and determinants of antimicrobial resistance. *Sci Rep* 2018;**8**:1–8.
- Baker S, The HC. Recent insights into *Shigella*: a major contributor to the global diarrhoeal disease burden. *Curr Opin Infect Dis* 2018;**31**:449.
- Belotserkovsky I, Sansonetti PJ. *Shigella* and enteroinvasive *Escherichia coli*. *Curr Top Microbiol Immunol* 2018;**416**:1–26.
- Dallman TJ, Chattaway MA, Mook P *et al.* Use of whole-genome sequencing for the public health surveillance of *Shigella sonnei* in England and Wales, 2015. *J Med Microbiol* 2016;**65**:882–4.
- Hawkey J, Paranagama K, Baker KS *et al.* Global population structure and genotyping framework for genomic surveillance of the major dysentery pathogen, *Shigella sonnei*. *Nat Commun* 2021;**12**:1–12.
- Holt KE, Baker S, Weill F-X *et al.* *Shigella sonnei* genome sequencing and phylogenetic analysis indicate recent global dissemination from Europe. *Nat Genet* 2012;**44**:1056–9.
- Hosangadi D, Smith PG, Kaslow DC *et al.* WHO consultation on ETEC and *Shigella* burden of disease, Geneva, 6–7th April 2017: meeting report. *Vaccine* 2019;**37**:7381–90.
- Mazi W, Sangal V, Saeed A *et al.* Rapid genotyping of *Shigella sonnei* by use of multiplex high-resolution melting. *J Clin Microbiol* 2015;**53**:2389–91.
- Pakbin B, Akhondzadeh Basti A, Khanjari A *et al.* Differentiation of *stx1A* gene for detection of *Escherichia coli* serotype O157: H7 and *Shigella dysenteriae* type 1 in food samples using high resolution melting curve analysis. *Food Sci Nutr* 2020a;**8**:3665–72.
- Pakbin B, Amani Z, Allahyari S *et al.* Genetic diversity and antibiotic resistance of *Shigella* spp. isolates from food products. *Food Sci Nutr* 2021a;**9**:6362–71.
- Pakbin B, Basti AA, Khanjari A *et al.* RAPD and ERIC-PCR coupled with HRM for species identification of non-dysenteriae *Shigella* species; as a potential alternative method. *BMC Res Notes* 2021b;**14**:1–6.
- Pakbin B, Didban A, Monfared YK *et al.* Antibiotic susceptibility and genetic relatedness of *Shigella* species isolated from food and human stool samples in Qazvin, Iran. *BMC Res Notes* 2021c;**14**:1–6.
- Pakbin B, Mahmoudi R, Mousavi S *et al.* Genotypic and antimicrobial resistance characterizations of *Cronobacter sakazakii* isolated from powdered milk infant formula: a comparison between domestic and imported products. *Food Sci Nutr* 2020b;**8**:6708–17.
- Phiri AF, Abia ALK, Amoako DG *et al.* Burden, antibiotic resistance, and clonality of *Shigella* spp. Implicated in community-acquired acute diarrhoea in Lilongwe, Malawi. *Trop Med Infect Dis* 2021;**6**:63.
- Ragupathi ND, Sethuvel DM, Inbanathan F *et al.* Accurate differentiation of *Escherichia coli* and *Shigella* serogroups: challenges and strategies. *New Microbes New Infect* 2018;**21**:58–62.
- Ranjbar R, Farahani A. *Shigella*: antibiotic-resistance mechanisms and new horizons for treatment. *Infect Drug Resist* 2019;**12**:3137.
- Reed GH, Kent JO, Wittwer CT. High-resolution DNA melting analysis for simple and efficient molecular diagnostics. *Pharmacogenomics* 2007;**8**:597–608.

- Rogawski McQuade ET, Shaheen F, Kabir F et al. Epidemiology of *Shigella* infections and diarrhea in the first two years of life using culture-independent diagnostics in 8 low-resource settings. *PLoS Negl Trop Dis* 2020;**14**:e0008536.
- Sangal V, Holt KE, Yuan J et al. Global phylogeny of *Shigella sonnei* strains from limited single nucleotide polymorphisms (SNPs) and development of a rapid and cost-effective SNP-typing scheme for strain identification by high-resolution melting analysis. *J Clin Microbiol* 2013;**51**:303–5.
- Shad AA, Shad WA. *Shigella sonnei*: virulence and antibiotic resistance. *Arch Microbiol* 2021;**203**:45–58.
- The HC, Bodhidatta L, Pham DT et al. Evolutionary histories and antimicrobial resistance in *Shigella flexneri* and *Shigella sonnei* in Southeast Asia. *Commun Biol* 2021;**4**:1–12.
- Torraca V, Holt K, Mostowy S. *Shigella sonnei*. *Trends Microbiol* 2020;**28**:696–7.
- Wittwer CT. High-resolution DNA melting analysis: advancements and limitations. *Hum Mutat* 2009;**30**:857–9.