

# Sprinting: a potential vaccine for hamstring injury?

Pascal Edouard<sup>1 2 3 4 5</sup>, Jurdan Mendiguchia<sup>6</sup>, Kenny Guex<sup>7</sup>, Johan Lahti<sup>8</sup>, Pierre Samozino<sup>9</sup>, and Jean-Benoît Morin<sup>8 10</sup>

<sup>1</sup>Inter-university Laboratory of Human Movement Science (LIBM EA 7424), University of Lyon, University Jean Monnet, F-42023. Saint Etienne, France, <sup>2</sup>Department of Clinical and Exercise Physiology, Sports Medicine Unit, University Hospital of Saint-Etienne, Faculty of Medicine, Saint-Etienne, France, <sup>3</sup>Medical Commission, French Athletics Federation (FFA), Paris, France, <sup>4</sup>Swiss Olympic Medical center, Centre de médecine du sport, Division de médecine physique et réadaptation, Centre Hospitalier Universitaire Vaudois, Lausanne, Switzerland, <sup>5</sup>European Athletics Medical & Anti Doping Commission, European Athletics Association (EAA), Lausanne, Switzerland, <sup>6</sup>Department of Physical Therapy, ZENTRUM Rehab and Performance Center, Barañain, Spain, <sup>7</sup>School of Health Sciences (HESAV), HES-SO University of Applied Sciences and Arts Western Switzerland, Lausanne, Switzerland, <sup>8</sup>Université Côte d'Azur, LAMHESS, Nice, France, <sup>9</sup>Univ Savoie Mont Blanc, Laboratoire Interuniversitaire de Biologie de la Motricité, EA 7424, F-73000 Chambéry, France, and <sup>10</sup>Sports Performance Research Institute New Zealand, School of Sport and Recreation, Auckland University of Technology, New Zealand

Hamstring | Sprint mechanics | Sports injury prevention | Track and field | Soccer

**G**ood athletes run fast. Running fast puts athletes at risk of hamstring muscle injuries (HMIs). Therefore, good athletes should avoid running fast, to avoid hamstring injuries. This apparently nonsensical syllogism is unfortunately the basis of a widespread avoidance of sprinting in many sports training environments.

Indeed, HMI is the main injury in sports requiring sprint acceleration, such as football(1) or athletics,(2) and thus a crucial issue for athletes and their supporting coaching and medical staffs. Due to the clear link between sprinting (i.e. sprinting constitutes the main HMI mechanism in those sports) and HMI occurrence,(3,4) sprinting is often blamed as the “foe”, which leads to HMI. Consequently, a widespread HMI prevention (and let us be honest, staff protection) measure, especially in training (games demands can hardly be altered), is to avoid sprint practice and monitoring up to maximal speed (≈20 to 60-m depending on the sport and level of practice). This approach is paradoxical since sprint acceleration performance can be considered one of the key performance indicators, justifying that sprinting should be the mean and the goal of any effective prevention strategy.

HMI prevention strategies have been developed and evaluated mainly based on isolated risk factors screening (strength evaluations)(5) or strengthening programmes without considering multiple risk factors and their interaction, and relationship with sprint action.(6) However, despite specific experimental contexts, real practice revealed that HMI still represents the most prevalent injury in sports involving sprinting.(1,2) This could be explained by the fact that (i) screening procedures are mainly focused on only one isolated risk factor (i.e. strength), (ii) hamstring muscle strength evaluations are mainly based on isolated, single joint, low-velocity, non-sprint-specific range of motion and torque evaluations,(7) and (iii) are not reliable enough in predicting HMIs,(5) (iv) prevention programmes were implemented at a group level without taking into account the individual characteristics and deficiencies, (v) showed low compliance,(6) and (vi) were mainly focused only on hamstring muscles strengthening using exercises which inevitably fall short in stimulating all the demands placed on the muscle group in sprinting cyclic.(7,8) These features of the current literature do not consider hamstring muscles as a constitutive element of the motor chain and the corresponding individual characteristics.(9) This emphasizes the need for a more global approach in HMI management (including prevention and rehabilitation),(9,10) in which we suggest that the complex and unique sprinting movement (leg interaction, elastic energy transfer, reflexes, kinematics, kinetics...) should be placed at the center of the approach.

In the risk of sounding redundant, sprinting is the only exercise that induces a sprint-specific hamstring muscles activation (both in terms of amount and timing): impossible to replicate by usual strengthening exercises that can reach only 18 to 75% of the electromyographic activity reached by hamstrings during sprinting.(8) It is thus irreplaceable as a component of a highly effective HMI rehabilitation programme, to reduce the risk of HMI re-injury in addition to improve sprint performance.(9) Contrary to rehabilitation programs only including strength and lengthening exercises, sprinting very likely induced tissue healing and repair, and helped athletes return to their sports including sprinting activities, in a combined “repair and prepare” strategy.(9) In addition, athletes who sprinted at high intensity (>95% of their maximal velocity) during sport practice showed a lower risk of lower limb injuries than those who produced lower maximal velocity (<85%).(11) Exposures to this activity was also important since an optimal maximal running velocity exposure was associated with a lower risk of lower limb injury,(11) and an acute increase in sprinting load was associated with large increased risk of HMI.(4) Therefore, adequate exposure to sprinting may be an effective “vaccine” against HMI. Like for a vaccine, the activity must be task-specific and should be adapted to the tissue and athletes’ capacity, in order to promote positive adaptation and a protective effect against the task-related risk, while having no side effect (such as HMI). Sprinting should be viewed by athletes and coaches as a mithridatization process.

Finally, we think that sprinting should be considered in the HMI prevention process, not only as a part of the problem but also, and more importantly, as a part of the solution. For the reason detailed above, we think that sprinting is irreplaceable as one of the HMI prevention measures (in primary and secondary prevention) in a global approach.(7,9,10) We also think that sprint performance improvement and HMI prevention are the same “fight”, with quite similar associated strategies. We think that sprint-oriented HMI prevention strategies can decrease injuries while also playing a “win-win” role in improving sprint performance. This approach could have a higher impact on athletes and coaches’ agreement, leading to likely higher compliance than methods previously proposed.(6) In this context, technical and medical staffs collaboration in a win-win performance-prevention strategy is highly relevant, since HMI affects all sports and stakeholders.

## References

1. Ekstrand J, Waldén M, Hägglund M. Hamstring injuries have increased by 4% annually in men’s professional football, since 2001: a 13-year longitudinal analysis of the UEFA

- Elite Club injury study. *Br J Sports Med* 2016;50:731–7. doi:10.1136/bjsports-2015-095359
2. Edouard P, Branco P, Alonso J-M. Muscle injury is the principal injury type and hamstring muscle injury is the first injury diagnosis during top-level international athletics championships between 2007 and 2015. *Br J Sports Med* 2016;50:619–30. doi:10.1136/bjsports-2015-095559
  3. Woods C, Hawkins RD, Maltby S, et al. The Football Association Medical Research Programme: an audit of injuries in professional football—analysis of hamstring injuries. *Br J Sport Med* 2004;38:36–41. doi:10.1136/bjism.2002.002352
  4. Duhig S, Shield AJ, Opar D, et al. Effect of high-speed running on hamstring strain injury risk. *Br J Sports Med* 2016;50:1536–40. doi:10.1136/bjsports-2015-095679
  5. Green B, Bourne MN, Pizzari T. Isokinetic strength assessment offers limited predictive validity for detecting risk of future hamstring strain in sport: a systematic review and meta-analysis. *Br J Sports Med* 2018;52:329–36. doi:10.1136/bjsports-2017-098101
  6. Goode AP, Reiman MP, Harris L, et al. Eccentric training for prevention of hamstring injuries may depend on intervention compliance: A systematic review and meta-analysis. *Br J Sports Med* 2015;49:349–56. doi:10.1136/bjsports-2014-093466
  7. Guex K, Millet GP. Conceptual framework for strengthening exercises to prevent hamstring strains. *Sport Med* 2013;43:1207–15. doi:10.1007/s40279-013-0097-y
  8. van den Tillaar R, Solheim JAB, Bencke J. Comparison of Hamstring Muscle Activation During High-Speed Running and Various Hamstring Strengthening Exercises. *Int J Sports Phys Ther* 2017;12:718–27. doi:10.16603/ijsp20170718
  9. Mendiguchia J, Martinez-Ruiz E, Edouard P, et al. A Multifactorial, Criteria-based Progressive Algorithm for Hamstring Injury Treatment. *Med Sci Sports Exerc* 2017;49:1482–92. doi:10.1249/MSS.0000000000001241
  10. Oakley AJ, Jennings J, Bishop CJ. Holistic hamstring health: not just the Nordic hamstring exercise. *Br J Sports Med* 2018;52:816–7. doi:10.1136/bjsports-2016-097137
  11. Malone S, Roe M, Doran DA, et al. High chronic training loads and exposure to bouts of maximal velocity running reduce injury risk in elite Gaelic football. *J Sci Med Sport* 2017;20:250–4. doi:10.1016/j.jsams.2016.08.005

**Copyright:** The articles published on *Science Performance and Science Reports* are distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.