Folia Forestalia Polonica, Series A – Forestry, 2016, Vol. 58 (2), 96–102

SHORT COMMUNICATION

DOI: 10.1515/ffp-2016-0011

First report of Geosmithia langdonii and Geosmithia spp. isolated from a decaying elm (Ulmus minor) in Geneva, Switzerland

Martine Hänzi, Bastien Cochard, Romain Chablais, Julien Crovadore, François Lefort 🖂

University of Applied Sciences and Arts Western Switzerland, Geneva Institute for Technology Architecture and Landscape, Institute Land Nature and Environment, Plants and Pathogens Group, 50 route de Presinge, 1254 Jussy, Switzerland, e-mail: francois.lefort@hesge.ch

Abstract

The mortality of a young elm *Ulmus minor* in 2014 in Geneva prompted a search for the microorganisms potentially involved. Symptoms included foliar chlorosis and wilting followed by defoliation of branches. Wood symptoms included a brown streaking of sap wood and brown stains in trunk and branches. The comparison of the resulting ITS rDNA sequences to the NCBI Nucleotide database allowed to identify 10 different organisms. The genus *Geosmithia* represented 48% of the isolates belonging to three species: *Geosmithia langdonii* (7 isolates) and 2 unknown morphologically and genetically different *Geosmithia* sp. 1 and sp. 2 (4 isolates). *Geosmithia* species are very little known ascomycetes, which have been recently shown to be opportunistic pathogens on broadleaved trees and conifers, living as saprobes in galleries of many bark beetle species. In the case described here, *Geosmithia langdonii*, and the unknown *Geosmithia* species were found in symptomatic wood while bark beetle galleries were found in close regions of the symptomatic wood. *Geosmithia langdonii* was the major fungus retrieved from the symptomatic wood and could have contributed, along with other identified fungal species, to a pathogenic complex producing symptoms similar to the ones of the Dutch Elm Disease and led to the dieback of this elm tree. *Geosmithia langdonii* and 2 yet unknown *Geosmithia* species (sp. 1 and sp. 2), different from any other reported *Geosmithia* species are reported from an elm tree in Switzerland for the first time.

Key words

Dutch elm disease Geosmithia langdonii, Geosmithia spp., elm, Ulmus minor, endophytes

Dutch elm disease (DED) caused by *Ophiostoma ulmi* and *O. novo-ulmi* is a matter of concern for city corporations, using elm trees in urban plantations. The death occurrence of a young elm in Sept. 2014, in Geneva (lat. 46.1984, long. 6.1423) in an urban plantation of several *Ulmus minor* trees, after a few months of decay

symptoms, which appeared end of May 2014 (Fig. 1), prompted a search for the microorganisms potentially involved. Symptoms included foliar chlorosis and wilting followed by defoliation of branches. No symptoms were observed on the other elms trees of this plantation. The tree was felled 2 weeks after death. A large



Figure 1. Wilting symptoms on an elm tree on 20th June 2014

branch and a small branch were cut in sections. One section of each was surface disinfected with ethanol 70% and sampled by coring, with a Pressler hand auger, in symptomatic regions as shown on Figure 2A. Wood symptoms included a brown streaking of sap wood and brown stains in trunk and branches. Three wood cores (5 mm diam.) were cultivated in Petri dishes on Potato Glucose Agar (PGA) medium (Roth AG, Switzerland) plus ampicillin (50 μ g/mL) (Roth AG) at room temperature (Fig. 3) and 23 fungal organisms were isolated in pure cultures (3 isolates from the trunk section sample, 19 isolates from the large branch section and 1 from the small branch section) on PGA medium. All isolates were observed and classified morphologically with the help of a dissecting microscope.

Subcultures for each morphologically different organism were grown in 50 mL Potato Glucose Broth (PGB) medium (Roth AG) plus ampicillin (50 μ g/mL) in 150 mL sterile Erlenmeyer flasks, under shaking (100 rpm) at room temperature for one week. Fungal DNA of 14 selected isolates was extracted following

a modified protocol adapted from Lefort and Douglas (1999) and assayed in a Nanodrop ND-1000 spectrophotometer (Wilmington, DE, USA) in order to adjust the final DNA concentration to 50 ng/ μ l in ultrapure water. PCR of the internal transcribed spacer rDNA (ITS1-5.8S-ITS2) was carried out with the primers pair ITS 4 and ITS 5 (White et al 1990). PCR reactions were performed in a Thermocycler Biometra (Göttingen, Germany) using the Taq polymerase and buffer from Biotools (Madrid, Spain) according to the following programme: initial denaturation for 3 min at 95°C, 38 cycles of 60 s at 95°C, 45 s at 56°C, 60 s at 72°C terminated by a final elongation of 6 min at 72°C. Reactions were carried out in 50 µL final volume at a final concentration of 1.5 mM MgCl₂, 100 µM each primer and 0.8 mM dNTPs. After amplification, PCR products quality was assessed by gel electrophoresis in a 1% agarose gel in 1x TBE (Tris



Figure 2. Symptomatic regions sampled on the trunk and branch sections



Figure 3. Wood cores on PGA medium (A); Fungal growth on cores on PGA medium (B)

89 mM, boric acid 89 mM, EDTA 2 mM, pH 8) and revealed by GelRedTM visualized through a UV transilluminator. PCR products were purified with the Wizard SV Gel and PCR Clean-Up System (Promega Corporation, Madison, USA). The sequencing of PCR products was carried out at Microsynth AG facilities (Balgach, Switzerland) and sequenced finally edited with Finch TV v 1.4.0 (Geospiza Inc., Seattle, WA, USA).

The comparison of the 14 resulting ITS rDNA sequences to the NCBI Nucleotide database (National Center of Biotechnology Information, Bethesda, USA) allowed identifying 10 different organisms, for which sequences have been registered in NCBI under the accessions KP790123-KP790136. The diversity of identified fungi is shown in Table 1. A number of morphologically identical isolates was not sequenced but are reported in Table 1. The genus Geosmithia represented 48% of the isolates belonging to three species: Geosmithia langdonii (7 isolates) and 2 unknown morphologically and genetically different *Geosmithia* sp. 1 (2 isolates) and Geosmithia sp. 2 (2 isolates). All these isolates were retrieved from the brown sap wood regions in the small and large branches, but not from the trunk as shown on Table 1. The 3 Geosmithia species were present in the large branch and one *Geosmithia* sp. was present in the small branch. Three representative isolates, Geosmithia langdonii UASWS1324, Geosmithia sp. 1 UASWS1325 and Geosmithia sp. 2 UASWS1334, have been deposited in the collection DSMZ-Deutsche Sammlung von Mikroorganismen und Zellkulturen (Braunschweig, Germany) under the reference numbers DSM 100090, DSM 100091 and DSM 100360, respectively. The appearance of these three Geosmithia species is shown on Figure 4 and conidia on a conidiophore of G. langdonii are shown on Figure 5. Most noticeably, none of the two fungal species responsible for DED, Ophiostoma ulmi or Ophiostoma novo-ulmi were isolated from symptomatic wood samples. The other identified fungi were potential endophytes such as Cladosporium sphaerospermum, 1 unknown Cladosporium sp., 1 Aspergillus sp., Penicillium brevicompactum and 2 unknown Sordariomycetes sp. genetically close to Aureobasidium sp. or Coniochaeta sp. It is unclear if some of them could be part of a phytopathogenic complex in association with Geosmithia species, able to lead to the observed tree's decay.

Geosmithia species are very little known ascomycetes, which have been recently shown to be opportunistic pathogens on broadleaved trees and conifers, living as saprobes in galleries of many bark beetle species (Jankowiak et al. 2014; Kolařík et al. 2005, 2007). Unknown and undescribed *Geosmithia* species were already reported on elm (Kolařík et al. 2005, 2007). Among the 50 described *Geosmithia* species, nine have been found on elm (Kolařík et al. 2008). More recently Pepori et al. (2015) isolated 72 strains of 6 *Geosmithia* species associated with elms and elm beetles, 3 of these species remaining undescribed. One up to four different species could be retrieved from a single tree. This is

Sample origin	Sample type	Isolates codes	Isolates accessions	GenBank accessions	Isolate identity
A Trunk section	Core A1	A1.1.1	UASWS1321	KP790123	Cladosporium sphaerospermum
	Core A2	A2.1.1	UASWS1322	KP790124	Aspergillus sp.
	Core A3	A3.1.1	UASWS1323	KP790125	Penicillium brevicompactum
B Large branch section	Core B1	B1.1.1	UASWS1324	KP790126	Geosmithia langdonii
		B1.1.2			Geosmithia langdonii
		B1.1.3	UASWS1325	KP790127	Geosmithia sp. 1
		B1.1.4			Geosmithia langdonii
		B1.2.1	UASWS1326	KP790128	Geosmithia langdonii
		B1.2.2	UASWS1327	KP790129	Geosmithia sp. 1
		B1.2.3a	UASWS1328	KP790130	Geosmithia langdonii
		B1.2.3b	UASWS1329	KP790131	Cladosporium sp.
		B1.2.3c			Geosmithia langdonii
		B1.2.4			Geosmithia sp. 2
		B1.2.5	UASWS1330	KP790132	Cladosporium sp.
		B1.2.6	UASWS1331	KP790133	Geosmithia langdonii
	Core B3	B3.1.1	UASWS1332	KP790134	Sordariomycetes sp. 1
		B3.1.2			Sordariomycetes sp. 1
		B3.1.3			Sordariomycetes sp. 1
		B3.1.4	UASWS1333	KP790135	Sordariomycetes sp. 2
		B3.2.1			Sordariomycetes sp. 2
		B3.2.2			Sordariomycetes sp. 2
		B3.2.3			Sordariomycetes sp. 2
C Small branch section	Core C2	C2.2.1	UASWS1334	KP790136	Geosmithia sp.

Table 1. Identification of isolated fungi from wood samples (sample origin, sample type, isolates codes, isolates accession in the collection of the Plants and Pathogens research group, GenBank accession and isolate identities)

therefore congruent with the observations of three different Geosmithia species reported here in a single tree. The comparison of the ITS DNA sequences of these 3 undescribed species, Geosmithia sp. 2, Geosmithia sp. 5 and Geosmithia sp. 20 (Pepori et al. 2015) to the species Geosmithia sp. 1 and Geosmithia sp. 2 of the present study yielded 91%, 91%, and 99% identity over 83% of the sequence length, respectively for Geosmithia sp. 1 93% identity, over 79% of the sequence length with each for *Geosmithia* sp. 2, suggesting that the diversity of Geosmithia species in elms is even greater. In the present case, galleries of bark beetles were observed in the wood of the trunk section (Fig. 2B) but not in the branches. An isolate of an undescribed Geosmithia sp. was found associated once with Dutch elm disease (Scala et al. 2007) in Italy. G. langdonii was also found

on elm in a survey of fungi associated to elms (Pepori 2012) and recently identified as a bark-beetle associate and an endophyte of *Ouercus agrifolia* in California (McPherson et al. 2013). Similarly to Ophiostoma ulmi (or O. novo-ulmi) in DED, the pathogenic Geosmithia species have most of the time been shown to be associated with bark beetle species. As stated by Kolařík et al. (2008), Geosmithia species would be dispersed by Phloeophagus bark beetles, with which the association might be very stable and specific, leading to symbiosis and speciation of Geosmithia species in some cases. That was the case for *Geosmithia morbida*, the agent of the Thousand Cankers Disease of walnut tree species in Europe and the USA (Kolařík et al. 2011; Zerillo et al. 2014; Montecchio et al. 2014), which has been shown to be associated with the bark beetle Pityophtorus juglan-



Figure 4. Relative proportions of fungal isolates



Figure 5. Colonies of *Geosmithia langdonii* isolate UASWS1324 v, *Geosmithia* sp. 1 isolate UASWS1325 (B) et *Geosmithia* sp. 2 isolate UASWS1334 (C) grown on PGA medium

dis. Geosmithia pallida was found associated with the bark beetle Pseudopityophthorus pubipennis in foamy bark canker in Quercus agrifolia in California (Lynch et al. 2014). Other species such as Geosmithia fassatiae, G. langdonii and G. obscura were discovered in association with Scolytus intricatus on several oak species (Kolařík et al. 2005, 2007). Geosmithia langdonii was also described as associated to Platypus cylindrus in the dieback of cork oaks (Quercus suber) in Algeria (Belhoucine et al. 2011). If pathogenicity of Geosmithia langdonii has not yet been proved, its effects on the growth of broadleaved trees and on roots have been demonstrated by Cízková et al. (2005). Geosmithia langdonii has also been often isolated in association with Ophiostoma ulmi and O. novo-ulmi in dying elms affected by DED (Pepori 2012; Bettini et al. 2014; Pepori et al. 2014). Besides these reported cases of plant pathogenicity, one Geosmithia species, Geosmithia argillacea has been shown to cause an invasive mycosis in humans associated to a cystic fibrosis (Giraud et al. 2010; De Ravin et al. 2011).

In the case described here, *Geosmithia langdonii*, and 2 other unknown *Geosmithia* species. *Geosmithia* sp. 1 and sp. 2 were found in symptomatic wood while bark beetle galleries are found in close regions of the symptomatic wood. *Geosmithia langdonii* was the major fungus retrieved from the symptomatic wood and could have taken part to a pathogenic complex contributing to DED symptoms and dieback of this elm tree. The hypothesis that this latter species, or the two unknown *Geosmithia* species, could not be part of a pathogenic complex cannot be completely withdrawn in absence of inoculations experiments confirming their pathogenicity. It cannot either be excluded that these species could be common endophytes of elm trees.

The main finding of this study is to report, for the first time, *Geosmithia langdonii* and 2 unknown species of *Geosmithia*, different from any known species, from an elm tree in Switzerland.

Acknowledgements

Mrss. Roger Beer and Bertrand Favre from the General Directorate of Agriculture and Nature of the Republic and Canton Geneva and the Geneva City Hall are acknowledged for funding these works.

REFERENCES

- Belhoucine L., Bouhraoua R.T., Meijer M., Equihua-Martinez A., Pujade-Villar J. 2011. Mycobiota associated with *Platypus cylindrus* (*Coleoptera: Curculionidae, Platypodidae*) in cork oak stands of North West Algeria, Africa. *African Journal of Microbiology Research*, 5 (25), 4411–4423.
- Bettini P.P., Frascella A., Kolařík M., Comparini C., Pepori A.L., Santini A., Scala F., Scala A. 2014. Widespread horizontal transfer of the cerato-ulmin gene between *Ophiostoma novo-ulmi* and *Geosmithia* species. *Fungal Biology*, 11, 663–674.
- Cízková D., Srůtka P., Kolarík M., Kubátová A., Pazoutová S. 2005. Assessing the pathogenic effect of *Fusarium, Geosmithia and Ophiostoma* fungi from broad-leaved trees. *Folia Microbiology*, 50, 59–62.
- De Ravin S.S., Challipalli M., Anderson V., Shea Y.R., Marciano B., Hilligoss D., Marquesen M., DeCastro R., Liu Y.-C., Sutton D.A., Wickes B.L., Kammeyer P.L., Sigler L., Sullivan K., Kang E.M., Malech H.L., Holland S.M., Zelazny A.M. 2011. *Geosmithia argillacea*: An emerging cause of invasive mycosis in human chronic granulomatous

disease. Clinical Infectious Diseases, 52 (6), e136e143.

- Giraud S., Pihet M., Razafimandimby B., Carrère J., Degand N., Mely M., Favennec L., Dannaoui E., Bouchara J.-P., Calenda A. 2010. *Geosmithia argillacea*: an emerging pathogen in patients with cystic fibrosis. *Journal of Clinical Microbiology*, 48 (7), 2381–2386.
- Jankowiak R., Kolařík M., Bilańskic P. 2014. Association of *Geosmithia* fungi (*Ascomycota: Hypocreales*) with pine and spruce-infesting bark beetles in Poland. *Fungal Ecology*, 11, 71–79.
- Kolařík M., Kubátová A., Cepicka I., Pazoutová S., Srůtka P. 2005. A complex of three new whitespored, sympatric, and host range limited *Geosmithia* species. *Mycological Research*, 109, 1323–1336.
- Kolařík M., Kostovcík M., Pazoutová S. 2007. Host range and diversity of the genus *Geosmithia (Ascomycota: Hypocreales)* living in association with bark beetles in the Mediterranean area. *Mycological Research*, 111, 1298–1310.
- Kolařík M., Kubátová A., Huler J., Pazoutová S. 2008. Geosmithia fungi are highly diverse and consistent bark beetle associates: evidence from their community structure in temperate Europe. *Microbial Ecol*ogy, 55, 65–80.
- Kolařík M., Freeland M., Utlet C., Tisserat N. 2011. Geosmithia morbida sp. nov., a new phytopathogenic species living in symbiosis with the walnut twig beetle (*Pityophthorus juglandis*) on Juglans in USA. Mycologia, 103, 325–332.
- Lefort F., Douglas G.C. 1999. An efficient micro-method of DNA isolation from mature leaves of four hardwood tree species *Acer, Fraxinus, Prunus* and *Quercus. Annals of Forest Science*, 56, 259–263.
- Lynch S.C., Wang D.H., Mayorquin J.S., Rugman-Jones P.F., Stouthamer R., Eskalen A. 2014. First Report of *Geosmithia pallida* causing foamy bark canker, a new disease on coast live oak (*Quercus agrifolia*), in association with *Pseudopityophthorus pubipennis* in California. *Plant Disease*, 98 (9), 1276–1276.
- McPherson B.A., Erbilgin N., Bonello P., Wood D.L. 2013. Fungal species assemblages associated with *Phytophthora ramorum* infected coast live oaks following bark and ambrosia beetle colonization in

northern California. *Forest Ecology and Management*, 291, 30–42.

- Montecchio L., Faccoli M. 2014. First record of Thousand Cankers disease *Geosmithia morbida* and walnut twig beetle *Pityophthorus juglandis* on *Juglans nigra* in Europe. *Plant Disease*, 98 (5), 696.
- Pepori A. 2012. Caratterizzazione e ruolo delle comunità di *Geosmithia* spp. su olmo. Ph.D. thesis, University of Florence, Florence, Italy.
- Pepori A., Luchi N., Kolarik M., Santini A. 2014. Role of fungi of the gen. *Geosmithia* in relation to the pathosystem elm-elm disease (DED). The IUFRO Incubator 2014, Abstract Volume, UFRO 2014 World Congress.
- Pepori A.L., Kolařík M., Bettini P.P., Vettraino A.M., Santini A. 2015. Morphological and molecular characterisation of *Geosmithia* species on European elms. *Fungal Biology*, 119 (11), 1063–1074.
- Scala A., Comparini C., Carresi L., Tegli S. 2005. The gene encoding cerato-ulmin, an *Ophiostoma*-produced protein involved in the Dutch elm disease, has been introgressed or horizontally transferred in an unrelated species of the genus *Geosmithia*. *Journal of Plant Pathology*, 87 (4), 305.
- Scala A., Comparini C., Tegli S., Scala F. 2007. A nonophiostoma fungus express the gene encoding the hydrophobin cerato-ulmin. *Journal of Plant Pathology*, 89 (2), 233–240.

- White T.J., Bruns T., Lee S., Taylor J. 1990. Amplification and direct sequencing of fungal ribosomal genes for phylogenetics. In: PCR protocols, a guide to methods and applications (eds.: M.A. Innis, D.H. Gelfand, J. Shinsky, T.J. White). Academic Press, San Diego.
- Zerillo M.M., Ibarra Caballero J., Woeste K., Graves A.D., Hartel C., Pscheidt J.W., Tonos J., Broders K., Cranshaw W., Seybold S.J., Tisserat N. 2014. Population structure of *Geosmithia morbida*, the causal agent of Thousand Cankers disease of walnut trees in the United States. *PLoS ONE*, 9 (11), e112847.
- http://www.ncbi.nlm.nih.gov/nuccore/814603658 Geosmithia langdonii isolate UASWS1324 18S ribosomal RNA gene, partial sequence; internal transcribed spacer 1 and 5.8S ribosomal RNA gene, complete sequence; and internal transcribed spacer 2, partial sequence.
- http://www.ncbi.nlm.nih.gov/nuccore/814603659 Geosmithia sp. 1 UASWS1325 18S ribosomal RNA gene, partial sequence; internal transcribed spacer 1 and 5.8S ribosomal RNA gene, complete sequence; and internal transcribed spacer 2, partial sequence
- http://www.ncbi.nlm.nih.gov/nuccore/814603668 Geosmithia sp. 2 UASWS1334 18S ribosomal RNA gene, partial sequence; internal transcribed spacer 1 and 5.8S ribosomal RNA gene, complete sequence; and internal transcribed spacer 2, partial sequence.