



BMJ Open Risk of 30-day hospital readmission associated with medical conditions and drug regimens of polymedicated, older inpatients discharged home: a registry-based cohort study

Filipa Pereira ^{1,2}, Henk Verloo ^{2,3}, Taushanov Zhivko,⁴ Saviana Di Giovanni,^{2,5} Carla Meyer-Masseti,⁶ Armin von Gunten,³ Maria Manuela Martins,^{1,7} Boris Wernli⁸

To cite: Pereira F, Verloo H, Zhivko T, *et al*. Risk of 30-day hospital readmission associated with medical conditions and drug regimens of polymedicated, older inpatients discharged home: a registry-based cohort study. *BMJ Open* 2021;**11**:e052755. doi:10.1136/bmjopen-2021-052755

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2021-052755>).

Received 25 April 2021
Accepted 01 July 2021



© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Filipa Pereira;
filipa.pereira@hevs.ch

ABSTRACT

Objectives The present study analysed 4 years of a hospital register (2015–2018) to determine the risk of 30-day hospital readmission associated with the medical conditions and drug regimens of polymedicated, older inpatients discharged home.

Design Registry-based cohort study.

Setting Valais Hospital—a public general hospital centre in the French-speaking part of Switzerland.

Participants We explored the electronic records of 20 422 inpatient stays by polymedicated, home-dwelling older adults held in the hospital's patient register. We identified 13 802 hospital readmissions involving 8878 separate patients over 64 years old.

Outcome measures Sociodemographic characteristics, medical conditions and drug regimen data associated with risk of readmission within 30 days of discharge.

Results The overall 30-day hospital readmission rate was 7.8%. Adjusted multivariate analyses revealed increased risk of hospital readmission for patients with longer hospital length of stay (OR=1.014 per additional day; 95% CI 1.006 to 1.021), impaired mobility (OR=1.218; 95% CI 1.039 to 1.427), multimorbidity (OR=1.419 per additional International Classification of Diseases, 10th Revision condition; 95% CI 1.282 to 1.572), tumorous disease (OR=2.538; 95% CI 2.089 to 3.082), polypharmacy (OR=1.043 per additional drug prescribed; 95% CI 1.028 to 1.058), and certain specific drugs, including antiemetics and antinauseants (OR=3.216 per additional drug unit taken; 95% CI 1.842 to 5.617), antihypertensives (OR=1.771; 95% CI 1.287 to 2.438), drugs for functional gastrointestinal disorders (OR=1.424; 95% CI 1.166 to 1.739), systemic hormonal preparations (OR=1.207; 95% CI 1.052 to 1.385) and vitamins (OR=1.201; 95% CI 1.049 to 1.374), as well as concurrent use of beta-blocking agents and drugs for acid-related disorders (OR=1.367; 95% CI 1.046 to 1.788).

Conclusions Thirty-day hospital readmission risk was associated with longer hospital length of stay, health disorders, polypharmacy and drug regimens. The drug regimen patterns increasing the risk of hospital readmission were very heterogeneous. Further research is needed to explore hospital readmissions caused solely by specific drugs and drug–drug interactions.

Strengths and limitations of this study

- The records of 20 422 hospitalisations involving 8878 different polymedicated, home-dwelling, older patients readmitted to hospital at least once were studied to determine the risk of 30-day hospital readmission.
- The study included 4-year data from a comprehensive hospital register (2015–2018).
- A whole series of sociodemographic and clinical parameters, medical conditions, and prescribed drugs were used to predict the probability of hospital readmission.
- Analyses were correlational and causality was not explored.
- Although the study considered statistical associations between drugs and hospital readmissions, it did not consider clinically diagnosed drug–drug interactions.

INTRODUCTION

Longitudinal studies have demonstrated that approximately 20% of home-dwelling older adults supported by home healthcare services experienced hospital readmission within 30 days of their discharge.^{1–3} For many older adults, readmission to an acute hospital is associated with a functional decline that has not always recovered by the time they are discharged.⁴ However, the systematic review by Hansen *et al*⁵ revealed wide-ranging estimates (5%–79%) of how many hospital readmissions were preventable. The period between hospital discharge and readmission has not always been clearly stated in the literature, ranging from 30 days to 3 years. However, 30 days is the most frequently used in public health policy when measuring healthcare system performance.^{6–8}



Numerous determinants have been identified and associated with hospital readmissions, for example, socio-demographic and individual characteristics, multimorbidity, and medical events.^{9 10} A substantial risk of 30-day hospital readmission has been associated with older inpatients treated for different diseases and surgical interventions involving hip fracture, cancer, bypass, acute cardiovascular events or complex surgery.¹¹ The reasons for hospital readmission after a surgical intervention are often not directly related to the surgery itself but rather to underlying chronic health conditions.¹² Thus, chronic diseases may play an important role in readmission risk, independent of the reason for the initial hospitalisation.^{13 14} Older adults' chronic diseases are not isolated health conditions; they can influence each other, and treatment for one disease may adversely affect another.¹⁵ For all these reasons, patterns of 30-day hospital readmissions may be very complex.¹⁶

Multimorbidity, in the case of two or more diseases,^{17 18} may require taking multiple medicines,¹⁹ known as polypharmacy (PP) when daily intake involves five or more drugs.²⁰ Increasing incidences of multimorbidity with age, and consequently PP, add to the complexity of managing older inpatients' drug prescriptions, particularly at hospital discharge.^{21 22} PP and inadequate drug management are significant risk factors for adverse drug events (ADEs)—the most common postdischarge complications—alongside hospital-acquired infections and procedural complications.^{23 24} ADEs resulting from inappropriate drug prescribing, discrepancies between prescribed and current regimens, poor adherence, and the inadequate surveillance of adverse effects frequently lead to hospital admissions, readmissions⁸ and other undesirable consequences such as increased morbidity, decreased autonomy, institutionalisation and even early death.^{25 26} A systematic review by El Morabet *et al*²⁷ indicated ADE rates of 18%–38% after hospital discharge and 4.5%–24% hospital readmission rates due to those ADEs. Because older adults use more drugs, they are at a greater risk of drug-related readmission. Numerous studies have found that nearly 30% of older inpatients experienced ADEs within 3 weeks of hospital discharge, almost three-quarters of which could have been prevented or lessened.^{10 28 29}

Despite the significant overall impact of ADEs on hospital readmission rates, little is known about the association of hospital readmission risk with medical conditions and drug regimens.^{30 31} El Morabet *et al* revealed the high prevalence of antibiotics, diuretics, vitamin K antagonists, opioids, antidiabetics, anticancer drugs, antihypertensives, digitalis glycosides, corticosteroids and psychotropic drugs in drug-related hospital readmissions.²⁷ Samoy *et al*³² reported that anticoagulants, hypoglycaemics, beta-blocking agents, antidepressants, calcium channel blockers and lenograstim were associated with high risk of hospital readmission. A retrospective patient record study by Teymoorian *et al*³³ reported that anticoagulants and antiplatelet agents, diuretics and

antihypertensives, and opioids were associated with a high risk of persons aged 80 years old or more being readmitted to hospital within 30 days. Blanc *et al* reported the readmission scores of different drugs in a large sample of 10374 adult hospital admissions in general medicine. Taking beta-blocking agents, calcium channel blockers, diuretics, hypoglycaemic drugs or opioids was a significant risk for 30-day readmission.⁹

Besides the higher risk of drug-related hospital readmission, some studies have also investigated the associations between combining drugs—a common practice when treating complex diseases or coexisting medical conditions—and drug-related hospital readmissions. Although using multiple drugs may be good clinical practice and compliant with guidelines for treating certain diseases, one significant consequence of combining drugs is that patients face much higher risk of ADEs, which can be caused by drug–drug interactions.^{34–36} ADEs can emerge because a drug's pharmacokinetics and pharmacodynamics change if taken with another drug.³⁶ Moura *et al*³⁷ found that participants with potential drug–drug interactions on their prescription list had 2.4 times higher adjusted OR of being readmitted.

Even though some studies have reported high numbers of readmissions among home-dwelling older patients for a variety of drugs,³⁸ this health issue was mostly investigated using prospective or cross-sectional studies with small samples. More insight is needed into the patterns of drug-related hospital readmissions and risk factors in order to design better interventions for addressing ADEs.^{39 40} As part of a broader project,⁴¹ the present study's goal was to use hospital register data to prioritise risk factors for hospital readmission. We hypothesised that sociodemographic characteristics, medical conditions (defined using the WHO's International Classification of Diseases, 10th Revision (ICD-10) and the Swiss Classification of Surgical Interventions (CHOP)) and drug prescriptions (based on the WHO's Anatomical Therapeutic Chemical (ATC) Classification System) were significant risk factors for 30-day hospital readmission for discharged older adults.

MATERIALS AND METHODS

Study design

This longitudinal study (2015–2018) used data on a population cohort taken from a hospital registry composed of 140 variables. These were used to investigate the associations between risk of 30-day hospital readmission and the medical conditions and drug regimens of polymedicated older inpatients discharged home. The study was performed with close regard to the RECORD (REporting of studies Conducted using Observational Routinely collected health Data) statement.⁴²

Population and data collection

Our custom, 4-year, registry-based data set included polymedicated inpatients (five or more drugs prescribed at

hospital discharge), aged 65 years old or more, living in their own homes and hospitalised at least once at the Valais Hospital (a public general hospital in the French-speaking part of Switzerland). This specific population was selected due to its increased risk of hospital readmission.^{10 28 29} Older inpatients hospitalised once only or who died during hospitalisation were excluded, as were those hospitalised for fewer than 24 hours (the criterion to count as 'hospitalised' in Switzerland). Valais Hospital's register contains a comprehensive electronic health record composed of 140 variables routinely collected during hospital stays. However, no electronic patient records were available for adult psychiatry for 2015–2018. The extracted patient data contained sociodemographic characteristics, medical and surgical diagnoses, and routinely assessed clinical data (such as gait, falls risk or hearing) from hospitalised patients with at least five prescribed drugs at discharge. Medical and surgical diagnoses were coded based on the ICD-10 and CHOP.⁴³ Drug classification was based on the WHO's ATC Classification System.⁴⁴

The strategy for transforming and synthesising the data extracted from the register's multiple data set sources was based on Olsen's register-based methodological considerations⁴⁵ and has been documented elsewhere.⁴⁶ Our data set was composed of 20 422 hospital admission records running from January 2015 to December 2018, with similar numbers of annual hospital admissions: 5134, 5095, 5125 and 5068, respectively.

Data set customisation for predictive analysis

The data set was recoded and customised to identify the frequency of older patients' hospital admissions. Each subject's unique identifier was used to distinguish their different hospital stays from 2015 to 2018. The data set included 13 802 readmissions involving 8878 different older inpatients discharged home, readmitted to hospital within 30 days and whose data were complete (no missing values).

Sociodemographic and clinical data were considered independent variables and used to compute the predictive models. Readmission following discharge home was defined as the dependent variable of interest and was dichotomised (0=no, 1=yes) based on 30-day readmission between 2015 and 2018. Furthermore, the custom data set was composed of six clinical clusters based on agglomerative hierarchical clustering methods for identifying clinically relevant characteristics and representing older inpatients' health status. Medical status and drugs data were recoded and copied to an exploitable population database.⁴⁶

Sociodemographic variables and length of stay

The sociodemographic data set—almost exclusively composed of ordinal variables—included two categorical variables (sex and place of discharge from hospital) and three continuous variables (age and admission and discharge dates). Sex and age were included in

the analysis as sociodemographic control variables. Age was considered a continuous variable as its progressive impact has been proven in preliminary investigations and previous studies.⁴⁷

Health variables

Numerous variables were used to describe older patients' health status during each hospital stay. The health data set was composed of 23 categorical variables: 21 measured as ordinal variables (mobility, changing position, falls in the last year, etc) and two measured as nominal variables (altered gait and chronic pain). A cleaner, better-structured data set—composed of hierarchical clusters—was obtained in a previous study combining empirical and best-practice statistical approaches.⁴⁶ Three of six preliminarily computed hierarchical clusters were included in the modelling analysis as confounding variables: the mobility cluster, the dependency on the activities of daily living cluster and the mental state cluster.⁴⁶ These three clusters were selected because of their significant contributions to hospital readmissions.^{48–50} The data set of medical information was composed of patients' principal medical diagnosis and four secondary medical diagnoses, based on ICD-10. Finally, the year of hospitalisation was introduced as a control variable, based on the fact that earlier admission to hospital during this period led to a higher probability of unplanned readmissions during the entire period covered.

Included drugs

The hospital data set showed that discharged patients had been prescribed 2370 different drugs. Drug prescriptions were considered continuous, classified according to the WHO's ATC Classification System⁵¹ and then included in the predictive model as independent variables. To ensure robust statistical results, the model only included drug categories prescribed to at least 30 inpatients who were readmitted within 30 days. Online supplemental file 1 presents the prescribed ATC classified drugs included in the predictive model as independent variables.

For statistical purposes, drug–drug interactions between different ATC drug classes⁵¹ were operationalised as dichotomised variables (0=no simultaneous use of drugs from both classes, 1=simultaneous use of drugs from both classes) and added to the previous model. Drug class interactions were selected based on a literature review, significant ORs and expert opinions.⁵²

Data analysis strategy

Data were extracted into a Microsoft Excel spreadsheet (Microsoft, Redmond, Washington, USA) and then imported into SPSS V.26.0 software. We examined statistical associations between hospital readmissions and patient age and sex, length of stay (LOS), principal and related ICD-10 diagnoses, CHOP interventions, and drug prescriptions during hospitalisations. A causality analysis between those variables was impossible given our retrospective data collection method, our inability



to calculate the time between drug intake and readmission, and the potential drug changes between hospitalisation sequences. We conducted a bivariate analysis relating the independent variables to 30-day readmission after discharge home from 2015 to 2018. Next, we calculated a series of multilevel logistic regression models for binary outcomes explaining the readmissions, within 30 days, of patients discharged home (0=no, 1=yes). These hierarchical models included two levels: the first level concerned hospital stays themselves, nested in the second level, that of individuals. First, we computed a baseline multilevel binary logistic regression model to estimate how sets of predictors influenced the probability of 30-day hospital readmission, which included individuals' characteristics, health conditions and hospital LOS. Second, we completed this baseline model with the drugs prescribed to older inpatients on their discharge home. Finally, to the baseline model completed with prescribed drugs, we added the known drug–drug interactions between different ATC drug classes, based on a literature review and expert opinions. The model computed each predictor's impact, other things being equal, by estimating its net impact, controlling for other factors (adjusted ORs). The model also considered correlations between each subject's different variables, which were generally not independent.⁵³ The model's random intercept design allowed each individual's intercept to vary, assuming that some unmeasured traits remained stable over time and allowing a better estimation of the model's parameters. The estimated parameters, on the other hand, had the same effect on every subject. Since the data were based on the whole population—not a sample—of polymedicated older inpatients discharged home from the Valais Hospital, the ORs' CIs and statistical tests were used to indicate the robustness of the relationships (they usually only make sense for statistical inference).

Patient and public involvement

Patients were not involved in the development of the research questions, study design, outcome measures and conduct of the study.

RESULTS

Descriptive results

The electronic records of 20 422 inpatient stays by polymedicated, home-dwelling older adults included the 13802 hospital readmissions of 8878 different older inpatients previously discharged home—an average of 1.55 inpatient hospital readmissions. The total sample's mean age was 77.77 years old (SD=7.48) and 57% were men (table 1). The average hospital LOS was 8.44 days (SD=7.58). At discharge, 25% of the sample had impaired mobility, 4% were impaired in their activities of daily living and 4% showed mental impairment. Our sample population averaged 4.58 (SD=0.92) ICD-10 diagnoses and 1.83 (SD=1.76) surgical interventions (CHOP) performed during hospitalisation. The selected medical diagnoses

Table 1 Sociodemographic and hospitalisation data for inpatient stays by polymedicated, home-dwelling adults aged 65 or more (N=13802)

Variables	Inpatient stays by polymedicated, home-dwelling adults aged 65 or more
Sex	
Stays by men (%)	7834 (56.8)
Stays by women (%)	5968 (43.2)
Age at discharge (years)	
Mean inpatient age at discharge (SD)	77.77 (7.48)
Minimum–maximum	65–106
Median (IQR 25–75)	77.00 (68.00–80.00)
65–69 (%)	2226 (16.1)
70–79 (%)	5811 (42.1)
80–89 (%)	4845 (35.1)
90 and more (%)	920 (6.7)
Year of discharge	
2015 (%)	3501 (25.4)
2016 (%)	3318 (24.0)
2017 (%)	3530 (25.6)
2018 (%)	3453 (25.0)
Length of stay (days)	
Mean (SD)	8.44 (7.58)
Minimum–maximum	1–149
Median (IQR 25–75)	7 (4–11)
Number of ICD-10 conditions	
Mean (SD)	4.58 (0.92)
Minimum–maximum	1–5
Median (IQR 25–75)	5 (5–5)
Principal ICD-10 diagnosis	
Circulatory (%)	3336 (24.2)
Infectious (%)	404 (2.9)
Respiratory (%)	1444 (10.5)
Trauma (%)	1043 (7.6)
Tumours (%)	1505 (10.9)
Number of CHOP surgical procedures	
Mean (SD)	1.83 (1.76)
Minimum–maximum	0–5
Median (IQR 25–75)	1 (0–3)
Number of medicines prescribed at hospital discharge	
Mean (SD)	8.95 (3.24)
Minimum–maximum	5–30
Median (IQR 25–75)	8 (7.50–16.00)

CHOP, Swiss Classification of Surgical Interventions; ICD-10, International Classification of Diseases, 10th Revision.

distinguished patients affected by circulatory (24%), infectious (3%) and respiratory (11%) diseases, as well as trauma (8%) and tumours (11%). On average, 8.95

(SD=3.24) drugs were prescribed per patient at hospital discharge.

Associations between 30-day hospital readmission risk and sociodemographic characteristics and medical conditions

The rate of 30-day hospital readmission for older patients discharged home was 7.8%. Bivariate associations with χ^2 tests showed significant differences between older inpatients' sociodemographic characteristics and medical conditions (table 2). Men showed a slightly higher proportion of 30-day hospital readmissions than women (8.2% vs 7.3%). However, age did not significantly affect the probability of 30-day readmission. More readmissions were also seen among older patients with a circulatory disease (8.2% vs 6.5%), those not affected by trauma (8.0% vs 5.8%) and especially those with a tumour (15.1% vs 6.9%). Multimorbidity also increased the risk of 30-day hospital readmissions—from 1.5% for older patients with a single ICD-10 condition to 8.8% for those with five.

Associations between 30-day hospital readmission risk and drugs

On average, older patients readmitted within 30 days had more prescribed drugs than those who were not readmitted (9.95 vs 8.87 drugs). We found a linear relationship between the 30-day readmission rate and the average number of prescribed drugs ($p>0.001$), which supported the absence of a cut-off point in this relationship (figure 1).

Among the most robust statistical associations (χ^2 tests) with 30-day hospital readmissions involved the classes of drugs including antineoplastics and immunomodulators (12.6% vs 7.6% for those not treated with them) and antiemetics and antinauseants (27.7% vs 7.7%). There was also a higher risk of 30-day hospital readmission among older inpatients taking drugs for functional gastrointestinal disorders (13.4% vs 7.4%) and antihypertensives (14.1% vs 7.7%) (table 3).

Baseline multivariate model

A baseline multivariate logistic regression model including older patients' sociodemographic and clinical variables, but not their prescribed drugs at discharge, was computed to predict 30-day hospital readmission after discharge home (table 4). Neither sex nor age had a significant impact. On the contrary, LOS had a significant impact (OR=1.014 for each additional day; 95% CI 1.006 to 1.021), as did mobility (OR=1.218 for older patients with an impaired mobility status; 95% CI 1.039 to 1.427). Dependence on the activities of daily living and mental health status showed no influence. Concerning diagnoses measured in the ICD-10, we found that older patients with a tumorous disease (OR=2.538; 95% CI 2.089 to 3.082) were much more susceptible to 30-day hospital readmission. Patients with circulatory pathologies showed no difference from the reference category (OR=0.938; 95% CI 0.783 to 1.124), nor did those with respiratory problems (OR=1.100; 95% CI 0.875 to 1.382), trauma

Table 2 30-day hospital readmission risk at different periods for different age groups (N=13 802 readmissions)

Variables	30-day hospital readmission (%)	P value
Complete sample	7.8	
Sex		*
Female vs male	7.3 vs 8.2	
Year-end age, in years		NS
65–69	7.5	
70–79	7.6	
80–89	8.4	
≥90	6.4	
Mobility cluster		NS
Preserved mobility vs impaired mobility	7.6 vs 8.5	
Activities in daily living (ADL)		NS
Full ADL ability vs impaired ADL	7.8 vs 7.2	
Cognitive status		NS
Preserved cognitive status vs cognitive impairment	7.8 vs 7.9	
ICD-10 diagnosis: circulatory problems		**
No vs yes	8.2 vs 6.5	
ICD-10 diagnosis: infection		NS
No vs yes	7.7 vs 9.9	
ICD-10 diagnosis: respiratory problems		NS
No vs yes	7.8 vs 8.0	
ICD-10 diagnosis: trauma		**
No vs yes	8.0 vs 5.8	
ICD-10 diagnosis: tumour		***
No vs yes	6.9 vs 15.1	
Number of ICD-10 conditions		***
1	1.5	
2	4.9	
3	3.6	
4	4.8	
5	8.8	
Number of surgical procedures (CHOP)		*
0	7.7	
1	7.8	
2	7.0	
3	7.3	
4	7.1	
5	9.7	
Year of discharge: 2015–2018		NS
2015	8.3	
2016	8.0	
2017	8.0	
2018	6.8	

* $P<0.05$, ** $P<0.01$, *** $P<0.001$.

CHOP, Swiss Classification of Surgical Interventions; ICD-10, International Classification of Diseases, 10th Revision; NS, non-significant.




Figure 1 Relationship between 30-day readmission rate and the number of prescribed drugs at discharge.

(OR=0.847; 95% CI 0.633 to 1.134) or infection-related problems (OR=1.381; 95% CI 0.964 to 1.977; $p=0.078$). Multimorbidity predicted a higher probability of readmission (OR=1.419 per additional ICD-10 condition; 95% CI 1.282 to 1.572), whereas the number of surgical procedures had no noticeable impact (OR=0.978; 95% CI 0.938 to 1.020). The year of hospital stay did have an impact, however, as the earlier the hospitalisation during the 4 years under review, the higher the probability of readmission (OR=0.933 per additional year; 95% CI 0.880 to 0.990).

Some variables that were non-significant in bivariate analyses became significant in multivariate analyses. This was because the results of multivariate analyses were controlled by all the other parameters and interpretations were made with 'other things being equal'. Also, the composition of subgroups could be very different in some bivariate analyses.

Prediction of 30-day hospital readmission and drug prescriptions

Table 5 presents the baseline logistic regression model completed with the drugs prescribed to older patients at discharge home that were significantly associated ($p \leq 0.05$) with 30-day hospital readmission. It was not possible to introduce the total number of drugs prescribed jointly in this model because of their collinearity with other drug variables. Non-significant drugs and other variables have been omitted from table 3 in order to simplify the presentation. The probabilities of 30-day hospital readmission are presented in descending order of discharged older patients' ORs for each additional unit of the drugs in question. Intake of antiemetics and antinauseants was very strongly linked to 30-day readmission (OR=3.216 for each additional drug unit taken; 95% CI 1.842 to 5.617), as were those of antihypertensives (OR=1.771; 95% CI 1.287 to 2.438), gastrointestinal drugs (OR=1.424; 95% CI 1.166 to 1.739), systemic hormonal preparations (OR=1.207; 95% CI 1.052 to 1.385) and vitamins (OR=1.201; 95% CI 1.049 to 1.374). On the contrary,

the intake of lipid-modifying agents was associated with a decrease in 30-day hospital readmissions (OR=0.841 for each drug from this class prescribed; 95% CI 0.732 to 0.967).

Drug interactions and 30-day hospital readmissions

The model considered drug class interactions for the (1) cardiovascular system*central nervous system, gastrointestinal system, and metabolism*cardiovascular system; (2) gastrointestinal system and metabolism*central nervous system; (3) cardiovascular system*anti-infectives; and (4) central nervous system*anti-infectives. The analysis was carried out controlling for the basic model's variables (table 4), and the table reports the ORs for each additional unit of the statistically significant drugs in question as well as for significant drug interactions. Antiemetics and antinauseants were very strongly associated with 30-day readmission (OR=3.222; 95% CI 1.844 to 5.630), as were drugs regulating the gastrointestinal tract (OR=1.428; 95% CI 1.169 to 1.744) and systemic hormones (OR=1.210; 95% CI 1.054 to 1.390). The joint intake of beta-blocking agents and drugs for acid-related disorders was significantly associated with 30-day hospital readmission (OR=1.367; 95% CI 1.046 to 1.788); this is the only significant drug interaction in table 4. On the contrary, lipid-modifying agents were associated with lower 30-day hospital readmission (OR=0.838), as were substances acting on the renin-angiotensin system (OR=0.892; 95% CI 0.796 to 0.999) (table 6).

DISCUSSION

The present study examined the records of 20 422 hospitalisations involving polymedicated home-dwelling older patients, eventually discharged home, to identify the risk of 30-day hospital readmission. These records were obtained from 4-year data of a comprehensive hospital register. The 8878 individual older patients readmitted to the Valais Hospital showed a 30-day hospital readmission rate of almost 8%, corroborating previously published all-cause hospital readmission rates among home-dwelling older patients.^{9 27} However, Jencks *et al*³ found a much higher 30-day readmission rate, reaching almost 20% among discharged older patients who had been hospitalised in acute medicine and surgery wards.³ As a bivariate association, multimorbid men were at a significantly higher risk of readmission than multimorbid women; however, in the adjusted multivariate analysis, this significance disappeared. Medical conditions, PP and multiple classes of prescribed drugs were all associated with higher 30-day readmission rates, in line with previous studies.^{27 54-56}

Our study found no significant differences in the risk of 30-day hospital readmission for men and women. However, some previous research found that men were more likely to forget to take their drugs or to not apply the changed drug dosages prescribed by their family physician, consequently increasing their risk of hospital readmission for

Table 3 30-day hospital readmissions for different classes of drugs based on ATC (N=13 802)

Drug class	30-day readmission with no drugs in this class (%)	30-day readmission with drugs in this class (%)	P value
First level, anatomical main group			
Blood and blood-forming organ drugs (B)	7.1	8.0	NS
Dermatologicals (D)	7.7	9.4	NS
Genitourinary system and sex hormones (G)	7.7	8.3	NS
Systemic hormonal preparations, excluding sex hormones and insulins (H)	7.4	9.5	***
Anti-infectives for systemic use (J)	8.0	7.2	NS
Antineoplastic and immunomodulating agents (L)	7.6	12.6	***
Drugs for the musculoskeletal system (M)	8.0	6.5	*
Antiparasitic products, insecticides and repellents (P)	7.8	6.6	***
Respiratory system drugs (R)	7.4	9.9	***
Sensory organ drugs (S)	7.8	8.4	NS
Second level, therapeutic subgroup			
Stomatological preparations (A01)	7.8	12.2	NS
Drugs for acid-related disorders (A02)	7.0	8.5	***
Drugs for functional gastrointestinal disorders (A03)	7.4	13.4	***
Antiemetics and anti-nauseants (A04)	7.7	27.7	***
Bile and liver therapy drugs (A05)	7.8	14.3	NS
Drugs for constipation (A06)	7.3	10.8	***
Antidiarrhoeals, intestinal anti-inflammatory/anti-infective agents (A07)	7.7	12.9	***
Digestives, including enzymes (A09)	7.8	10.0	NS
Drugs used in diabetes (A10)	7.4	9.5	***
Vitamins (A11)	7.5	9.9	***
Mineral supplements (A12)	7.4	8.8	**
Other alimentary tract and metabolism products (A16)	7.8	6.3	NS
Cardiac therapy (C01)	7.6	8.9	NS
Antihypertensives (C02)	7.7	14.1	***
Diuretics (C03)	7.2	9.8	***
Peripheral vasodilators (C04)	7.8	15.2	NS
Vasoprotective drugs (C05)	7.8	9.8	NS
Beta-blocking agents (C07)	7.1	8.6	***
Calcium channel blockers (C08)	7.7	8.6	NS
Agents acting on the renin-angiotensin system (C09)	8.7	7.1	***
Lipid-modifying agents (C10)	8.3	7.1	**
Anaesthetics (N01)	7.8	18.8	*
Analgesics (N02)	7.8	7.8	NS
Antiepileptics (N03)	7.7	9.0	NS
Drugs for Parkinson's disease (N04)	7.8	6.9	NS
Psycholeptics (N05)	6.8	9.3	***
Psychoanaleptics (N06)	7.8	7.7	NS
Other nervous system drugs (N07)	7.9	5.1	NS

*P<0.05, **P<0.01, ***P<0.001.

ATC, Anatomical Therapeutic Chemical Classification System; NS, non-significant.

drug-related problems.⁵⁷ Opposite results were found in a population-based study by Manteuffel *et al*,⁵⁸ with women being less likely than men to properly adhere to their drug prescriptions. These differences may indicate a need for

more personalised drug prescription and drug management to improve clinical outcomes. Further research should explore associations between different types of drugs and sex,^{58 59} but this topic was beyond the scope

**Table 4** Baseline multilevel logistic regression model using 30-day readmission (0=no; 1=yes) as the dependent variable associated with independent sociodemographic, LOS and clinical variables (N=13 802 observations for 8878 different inpatients readmitted to hospital)

Variables	OR*	P value >z	95% CI
Sex	1.079	0.285	0.938 to 1.242
Year-end age, in years	0.999	0.878	0.990 to 1.009
Hospital LOS, in days	1.014	0.000	1.006 to 1.021
Mobility cluster†	1.218	0.015	1.039 to 1.427
Dependency on the activities of daily living‡	0.794	0.248	0.537 to 1.174
Mental health status†	0.992	0.966	0.687 to 1.433
ICD 1st diagnosis: circulatory problems‡	0.938	0.491	0.783 to 1.124
ICD 1st diagnosis: infection‡	1.381	0.078	0.964 to 1.977
ICD 1st diagnosis: respiratory problems‡	1.100	0.414	0.875 to 1.382
ICD 1st diagnosis: trauma‡	0.847	0.265	0.633 to 1.134
ICD 1st diagnosis: tumour‡	2.538	0.000	2.089 to 3.082
Number of ICD	1.419	0.000	1.282 to 1.572
Number of CHOP	0.978	0.304	0.938 to 1.020
Number of drugs	1.043	0.000	1.028 to 1.058
Year: 2015–2018	0.933	0.022	0.880 to 0.990
Intercept	–	0.027	–

*Adjusted OR.

†0=good state; 1=impairment.

‡0=no; 1=yes.

CHOP, Swiss Classification of Surgical Interventions; LOS, length of stay.

of the present study. Another interesting issue regarding sex differences in hospital readmission rates is the study window. Some studies found higher rates among men than among women below 3-month readmissions. More extended time windows (eg, 1 year) revealed no significant sex differences.^{54 60} An analysis of our data set using a more extended readmission window might clarify this point and provide complementary knowledge about sex-associated hospital readmissions.

Our results indicated that ageing was not a risk factor for increased 30-day hospital readmission, in line with some previous publications.^{55 61} However, other research findings demonstrated that age was only positively associated with the likelihood of readmission up to 74 years old; above this, there no longer appeared to be any significant relationship between age and readmission.^{62 63} These contrasting results may be explained by the studies' designs, country settings, age of research populations or the medical conditions included.^{55 62 64}

Longer hospital stays were also associated with a higher risk of hospital readmission, in line with a cohort study

by Sud *et al*⁶⁵ concluding that an extended hospital LOS was associated with increased rates of all types of readmission, except for hospitalisation after heart failure, where a short LOS was associated with increased rates of readmission for cardiovascular disease and heart failure.

Our results indicated a significant positive association between the number of a patient's medical conditions and the 30-day hospital readmission rate, confirming other recent retrospective hospital register studies.^{66 67} More specifically, older patients with impaired mobility showed an increased risk of hospital readmission. This result was not surprising, bearing in mind that although these older patients were discharged home—and not to a nursing home—after their hospital stay, their health status might nevertheless require future readmission. Indeed, this corroborated publications about older patients discharged after orthopaedic treatment or who had been initially admitted for heart failure, myocardial infarction or pneumonia, but also presented with impaired mobility.^{68 69}

Cognitive impairment was not associated with increased 30-day hospital readmission rates, in line with findings from the systematic review by Pickens *et al*,⁷⁰ which pointed out that dementia had a modest impact on readmission rates. It was no surprise that inpatients hospitalised for cancer faced a high risk of readmission, corroborating prior studies by Burhenn *et al*, Chang *et al* and Butcher.^{71–73}

PP significantly increased the 30-day hospital readmission rate, but this result was based on the average number of drugs prescribed to the sample of readmitted patients versus those not readmitted. Although PP was confirmed as a strong determinant of 30-day hospital readmission in publications by Leendertse *et al*,^{74 75} our results showed a progressive linear relationship between PP and readmission rate, and this should be interpreted with caution. Despite our results and other publications and research underlining the challenge of PP among multimorbid older patients, there is no overall consensus about the best way to deal with the broad general relationship between PP and hospital readmission.⁷⁶

Our advanced statistical analysis demonstrated that some specific drugs and the concomitant use of specific drug combinations were significantly associated with 30-day readmission risk, although this was not unexpected and has been confirmed in previous publications.^{37 77} Mostly in line with the research findings of Zhang *et al*, drugs including hormones, antineoplastics, immunosuppressors, neoplastic antibiotics and bacterial vaccines were substantial risk factors for hospital readmission.⁷

In summary, extended hospital LOS, functional impairments, medical conditions and drugs have been demonstrated to be determinants of 30-day hospital readmission, although not all of them have clinically or pharmacologically relevant interpretations or explanations. Further research involving large samples is needed, notably to explore the drug–drug interactions with the highest risk of hospital readmissions. Statistical predictions of

Table 5 Multilevel logistic regression model results for the drugs prescribed to older patients at discharge home that had significant predictive values (OR) for 30-day hospital readmission (controlled for variables in the baseline model: table 4) (N=13 802 observations for 8878 different inpatients readmitted to hospital)

Variables	OR*	P value >z	95% CI
First level, anatomical main group			
Blood and blood-forming organs drugs (B)	1.089	0.041	1.003 to 1.181
Systemic hormonal preparations, excluding sex hormones and insulins (H)	1.207	0.007	1.052 to 1.385
Respiratory system drugs (R)	1.146	0.003	1.046 to 1.254
Second level, therapeutic subgroup			
Drugs for functional gastrointestinal disorders (A03)	1.424	0.001	1.166 to 1.739
Antiemetics and antinauseants (A04)	3.216	0.000	1.842 to 5.617
Drugs for constipation (A06)	1.195	0.018	1.031 to 1.386
Drugs used in diabetes (A10)	1.125	0.021	1.018 to 1.243
Vitamins (A11)	1.201	0.008	1.049 to 1.374
Antihypertensives (C02)	1.771	0.000	1.287 to 2.438
Diuretics (C03)	1.149	0.024	1.018 to 1.296
Beta-blocking agents (C07)	1.156	0.040	1.007 to 1.327
Lipid-modifying agents (C10)	0.841	0.015	0.732 to 0.967
Psycholeptics (N05)	1.130	0.009	1.031 to 1.238

*Adjusted OR.

potential drug–drug interactions provide important information for modelling drug combinations and identifying pairs of drugs whose combination creates an exaggerated response.⁹ As the association between the number of drugs and the risk of hospital readmission was linear, more advanced inferential statistics would be needed to clarify a cut-off point for the mean number of drugs that would significantly increase the readmission rate. In addition, problems involving adherence to prescriptions, social support networks, and stronger or weaker

primary healthcare structures can all influence hospital readmission rates.³⁹ According to some publications, nearly 70% of people aged over 65 make mistakes with their drugs.^{78 79} Information about drug adherence, drug underuse and overuse, drug changes and deprescription by family physicians, as well as medication management at home would contribute to a more comprehensive understanding of disease-related and drug-related 30-day hospital readmissions.

Table 6 Drugs and drugs interactions from ATC classes A and B with a significant risk of 30-day hospital readmission (controlled for variables in the baseline model: table 4) (N=13 802 observations for 8878 different inpatients readmitted to hospital)

Variables	OR*	P value >z	95% CI
First level, anatomical main group			
Blood and blood-forming organ drugs (B)	1.089	0.040	1.004 to 1.182
Systemic hormonal preparations, excluding sex hormones and insulins (H)	1.210	0.007	1.054 to 1.390
Respiratory system drugs (R)	1.149	0.003	1.049 to 1.258
Second level, therapeutic subgroup			
Antiemetics and antinauseants (A04)	3.222	0.000	1.844 to 5.630
Drugs for functional gastrointestinal disorders (A03)	1.428	0.000	1.169 to 1.744
Beta-blocking agents (C07) and drugs for acid-related disorders (A02)	1.367	0.022	1.046 to 1.788
Drugs for constipation (A06)	1.199	0.017	1.033 to 1.392
Agents acting on the renin–angiotensin system (C09)	0.892	0.049	0.796 to 0.999
Lipid-modifying agents (C10)	0.838	0.013	0.729 to 0.964

*Adjusted OR.

ATC, Anatomical Therapeutic Chemical Classification System.

Finally, it would be interesting to explore the risk of readmission according to different hospital wards. As psychiatric conditions are a frequent cause of rehospitalisation,⁸⁰ it would be relevant for future research to explore registries from adult psychiatry departments and investigate the hospital readmission risk faced by their inpatients.

Strengths and limitations

This study's main strength was its use of data recorded in a comprehensive register. We consider this retrospective study useful for clinical practice and future research because a whole series of sociodemographic and clinical parameters, medical conditions, and prescribed drugs were used to predict the probability of hospital readmission. Using both bivariate and multivariate analyses enabled an evaluation of the data's longitudinal nature.

Our study had several limitations, nevertheless. The design did not allow us to identify hospitalisations and readmissions lost to follow-up and to adjust our data for death outside the hospital. We were also unable to identify unnecessary hospitalisations or any bias towards hospitalisation rather than another healthcare solution for older inpatients. Our data set could not inform us about whether older inpatients had been first admitted to another hospital or were subsequently readmitted elsewhere during the study period. Because the reasons for hospital admission are not chosen from a list but are entered into the register as free descriptive text, these factors were not part of our data set, and the study was unable to explore the reasons for an admission's impact on 30-day rehospitalisation. Another limitation was the study's lack of formal screening methods to explain ADEs in detail, and it was impossible to distinguish between elective and urgent hospitalisations. Although the study considered statistical associations between drugs and rehospitalisations, it did not use clinically diagnosed drug–drug interactions. Finally, we were unable to consider any potential causality between PP and hospital readmission.

CONCLUSIONS

Hospital LOS, medical conditions, functional impairments and prescribed drugs were all critical factors in predicting hospital readmissions, thus affirming our hypotheses. Readmission patterns are complex and poorly understood because older patients often present with multiple chronic conditions, functional impairments and complex drug prescriptions. Hospital readmission is an underinvestigated topic deserving of additional, well-conducted, predictive research exploiting accurate longitudinal data from large samples.

Author affiliations

¹Institute of Biomedical Sciences Abel Salazar, University of Porto, Porto, Portugal

²School of Health Sciences, HES-SO University of Applied Sciences and Arts Western Switzerland, Sion, Switzerland

³Service of Old Age Psychiatry, Lausanne University Hospital, Lausanne, Switzerland

⁴Faculty of Psychology and Educational Sciences, University of Geneva, Geneva, Switzerland

⁵Pharmacy Benu Tavil-Chatton, Morges, Switzerland

⁶Institute for Primary Health Care, University of Bern, Bern, Switzerland

⁷Porto Higher School of Nursing, Porto, Portugal

⁸FORS, Swiss Centre of Expertise in the Social Sciences, University of Lausanne, Lausanne, Switzerland

Twitter Maria Manuela Martins @mmmartins1956

Acknowledgements The authors thank the partner hospital, including the hospital's data warehouse, for its valuable contributions.

Contributors BW, FP and HV had the original idea. BW, TZ, SDG, MMM and HV provided conceptual and methodological expertise to the study design and BW, FP, TZ, SDG, CM-M, AV-G and HV to data analysis and interpretation. BW, FP and HV were major contributors to writing the manuscript. All authors read, edited and approved the final manuscript.

Funding This study was supported by the Swiss National Science Foundation via grant number 407440_183434/1. This research was developed, in part, using grants from the Swiss National Science Foundation and the School of Health Sciences of the University of Applied Sciences and Arts Western Switzerland (HES-SO) Valais/Wallis.

Disclaimer The funders had no role in the design and conduct of the study, collection, management, analysis and interpretation of data, preparation, review or approval of the manuscript, or the decision to submit the manuscript for publication.

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval Ethical approval was obtained from the Human Research Ethics Committee of the Canton of Vaud (CER-VD, 2018-02196), thus permitting our partner hospital's data warehouse to provide the appropriate data set. Given the retrospective data source, obtaining consent from the patients concerned was impossible or posed disproportionate difficulties. The present study respects the legal requirements for research projects involving data reuse without consent, as set out in Art 34 from the Swiss Human Research Act (HTA).

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

ORCID iDs

Filipa Pereira <http://orcid.org/0000-0001-9207-4856>

Henk Verloo <http://orcid.org/0000-0002-5375-3255>

REFERENCES

- 1 Rayan-Gharra N, Shadmi E, Tadmor B, *et al*. Patients' ratings of the in-hospital discharge briefing and post-discharge primary care follow-up: the association with 30-day readmissions. *Patient Educ Couns* 2019;102:1513–9.
- 2 Kabue S, Greene J, Kipnis P, *et al*. The impact of Pharmacy-specific predictors on the performance of 30-day readmission risk prediction models. *Med Care* 2019;57:295–9.
- 3 Jencks SF, Williams MV, Coleman EA. Rehospitalizations among patients in the Medicare fee-for-service program. *N Engl J Med Overseas Ed* 2009;360:1418–28.

- 4 Covinsky KE, Palmer RM, Fortinsky RH, *et al.* Loss of independence in activities of daily living in older adults hospitalized with medical illnesses: increased vulnerability with age. *J Am Geriatr Soc* 2003;51:451–8.
- 5 Hansen LO, Young RS, Hinami K, *et al.* Interventions to reduce 30-day rehospitalization: a systematic review. *Ann Intern Med* 2011;155:520–8.
- 6 Davies EC, Green CF, Mottram DR, *et al.* Emergency re-admissions to hospital due to adverse drug reactions within 1 year of the index admission. *Br J Clin Pharmacol* 2010;70:749–55.
- 7 Zhang M, Holman C D'Arcy J, Price SD, *et al.* Comorbidity and repeat admission to hospital for adverse drug reactions in older adults: retrospective cohort study. *BMJ* 2009;338:a2752.
- 8 Bonnet-Zamponi D, d'Arailh L, Konrat C, *et al.* Drug-Related readmissions to medical units of older adults discharged from acute geriatric units: results of the optimization of medication in aged multicenter randomized controlled trial. *J Am Geriatr Soc* 2013;61:113–21.
- 9 Blanc A-L, Fumeaux T, Stirnemann J, *et al.* Development of a predictive score for potentially avoidable hospital readmissions for general internal medicine patients. *PLoS One* 2019;14:e0219348.
- 10 Stevenson JM, Davies JG, Martin F. Is medication related harm as a cause of readmission associated with the indicators of frailty? *Age and Ageing* 2018;47:ii19.
- 11 Brunner-La Rocca H-P, Peden CJ, Soong J, *et al.* Reasons for readmission after hospital discharge in patients with chronic diseases-Information from an international dataset. *PLoS One* 2020;15:e0233457.
- 12 Dharmarajan K, Hsieh AF, Lin Z, *et al.* Hospital readmission performance and patterns of readmission: retrospective cohort study of Medicare admissions. *BMJ* 2013;347:f6571.
- 13 Arora S, Patel P, Lahewala S, *et al.* Etiologies, trends, and predictors of 30-day readmission in patients with heart failure. *Am J Cardiol* 2017;119:760–9.
- 14 Shams I, Ajorlou S, Yang K. A predictive analytics approach to reducing 30-day avoidable readmissions among patients with heart failure, acute myocardial infarction, pneumonia, or COPD. *Health Care Manag Sci* 2015;18:19–34.
- 15 Krumholz HM, Wang K, Lin Z, *et al.* Hospital-Readmission risk – isolating Hospital effects from patient effects. *NEJM* 2017;377:1055–64.
- 16 Gruneir A, Fung K, Fischer HD, *et al.* Care setting and 30-day Hospital readmissions among older adults: a population-based cohort study. *CMAJ* 2018;190:E1124–33.
- 17 Marengoni A, Angleman S, Melis R, *et al.* Aging with multimorbidity: a systematic review of the literature. *Ageing Res Rev* 2011;10:430–9.
- 18 Valderas JM, Starfield B, Sibbald B, *et al.* Defining comorbidity: implications for understanding health and health services. *Ann Fam Med* 2009;7:357–63.
- 19 Maher RL, Hanlon J, Hajjar ER. Clinical consequences of polypharmacy in elderly. *Expert Opin Drug Saf* 2014;13:57–65.
- 20 Masnoon N, Shakib S, Kalisch-Ellett L, *et al.* What is polypharmacy? A systematic review of definitions. *BMC Geriatr* 2017;17:1–10.
- 21 Wastesson JW, Morin L, Tan ECK, *et al.* An update on the clinical consequences of polypharmacy in older adults: a narrative review. *Expert Opin Drug Saf* 2018;17:1185–96.
- 22 Rieckert A, Trampisch US, Klaaßen-Mielke R, *et al.* Polypharmacy in older patients with chronic diseases: a cross-sectional analysis of factors associated with excessive polypharmacy. *BMC Fam Pract* 2018;19:113.
- 23 Magill SS, Edwards JR, Bamberg W, *et al.* Multistate point-prevalence survey of health care-associated infections. *N Engl J Med* 2014;370:1198–208.
- 24 Coleman EA, Smith JD, Raha D, *et al.* Posthospital medication discrepancies: prevalence and contributing factors. *Arch Intern Med* 2005;165:1842–7.
- 25 Ferreri SP, Hughes TD, Snyder ME. Medication therapy management: current challenges. *Integr Pharm Res Pract* 2020;9:71–81.
- 26 Roux P, Verloo H, Santiago-Delefosse M. The spatial dimensions of medication management by home-dwelling older adults after hospital discharge. *Health Place* 2019;60:102230.
- 27 El Morabet N, Uitvlugt EB, van den Bemt BJJ, *et al.* Prevalence and preventability of drug-related Hospital readmissions: a systematic review. *J Am Geriatr Soc* 2018;66:602–8.
- 28 Forster AJ, Murff HJ, Peterson JF, *et al.* The incidence and severity of adverse events affecting patients after discharge from the hospital. *Ann Intern Med* 2003;138:161–7.
- 29 Yeo I, Cheung JW, Feldman DN, *et al.* Assessment of hospital readmission rates, risk factors, and causes after cardiac arrest: analysis of the US nationwide readmissions database. *JAMA Netw Open* 2019;2:e1912208–e.
- 30 Hauviller L, Eyvrard F, Garnault V, *et al.* Hospital re-admission associated with adverse drug reactions in patients over the age of 65 years. *Eur J Clin Pharmacol* 2016;72:631–9.
- 31 Davies EC, Green CF, Mottram DR, *et al.* Emergency re-admissions to hospital due to adverse drug reactions within 1 year of the index admission. *Br J Clin Pharmacol* 2010;70:749–55.
- 32 Samoy LJ, Zed PJ, Wilbur K, *et al.* Drug-Related hospitalizations in a tertiary care internal medicine service of a Canadian Hospital: a prospective study. *Pharmacotherapy* 2006;26:1578–86.
- 33 Teymoorian SS, Dutcher D, Woods M. Association between postdischarge adverse drug reactions and 30-day Hospital readmission in patients aged 80 and older. *J Am Geriatr Soc* 2011;59:948–9.
- 34 WHO. *Adherence to long-term therapies: evidence for action.* Geneva 2003.
- 35 Brahma DK, Wahlang JB, Marak MD, *et al.* Adverse drug reactions in the elderly. *J Pharmacol Pharmacother* 2013;4:91–4.
- 36 Davies EA, O'Mahony MS. Adverse drug reactions in special populations - the elderly. *Br J Clin Pharmacol* 2015;80:796–807.
- 37 Moura CSde, Tavares LS, Acurcio FdeA. [Hospital readmissions related to drug interactions: a retrospective study in a hospital setting]. *Rev Saude Publica* 2012;46:1082–9.
- 38 Spinks JM, Kalisch Ellett LM, Spurling G, *et al.* Adaptation of potentially preventable medication-related hospitalisation indicators for indigenous populations in Australia using a modified Delphi technique. *BMJ Open* 2019;9:e031369.
- 39 Rosen OZ, Fridman R, Rosen BT, *et al.* Medication adherence as a predictor of 30-day hospital readmissions. *Patient Prefer Adherence* 2017;11:801–10.
- 40 Pellegrin KL, Lee E, Uyeno R, *et al.* Potentially preventable medication-related hospitalizations: a clinical pharmacist approach to assessment, categorization, and quality improvement. *J Am Pharm Assoc* 2017;57:711–6.
- 41 Pereira F, Roux P, Santiago-Delefosse M, *et al.* Optimising medication management for polymedicated home-dwelling older adults with multiple chronic conditions: a mixed-methods study protocol. *BMJ Open* 2019;9:e030030.
- 42 Benchimol EI, Smeeth L, Guttman A, *et al.* The reporting of studies conducted using observational routinely-collected health data (record) statement. *PLoS Med* 2015;12:e1001885.
- 43 Surgical interventions classification S. *Swiss classification of surgical interventions (CHOP)*, 2016.
- 44 Parker S, Prince A, Thomas L, *et al.* Electronic, mobile and telehealth tools for vulnerable patients with chronic disease: a systematic review and realist synthesis. *BMJ Open* 2018;8:e019192.
- 45 Olsen J. Register-Based research: some methodological considerations. *Scand J Public Health* 2011;39:225–9.
- 46 Taushanov Z, Verloo H, Wernli B, *et al.* Transforming a patient registry into a customized data set for the advanced statistical analysis of health risk factors and for medication-related hospitalization research: retrospective hospital patient registry study. *JMIR Med Inform* 2021;9:e24205.
- 47 Andrade C. Age as a variable: continuous or categorical? *Indian J Psychiatry* 2017;59:524–5.
- 48 Gerka ALC, Pfingsthorn M, Eichelberg M, SCITEPRESS – Science and Technology Publications. A Clustering-based approach to determine a standardized statistic for daily activities of elderly living alone. *Proceedings of the 12th International Joint Conference on Biomedical Engineering Systems and Technologies*, Lda, 2019.
- 49 Trevithick L, Painter J, Keown P. Mental health clustering and diagnosis in psychiatric in-patients. *BJPsych Bull* 2015;39:119–23.
- 50 Delil S, Çelik RN, San S, *et al.* Clustering patient mobility patterns to assess effectiveness of health-service delivery. *BMC Health Serv Res* 2017;17:1–14.
- 51 WHO. The anatomical therapeutic chemical classification system with defined daily doses (ATC/DDD) Geneva: World Health organization, 2014. Available: <http://www.who.int/classifications/atcddd/en/> [Accessed 24 Mar 2020].
- 52 Haider SI, Johnell K, Thorslund M, *et al.* Trends in polypharmacy and potential drug-drug interactions across educational groups in elderly patients in Sweden for the period 1992 - 2002. *Int J Clin Pharmacol Ther* 2007;45:643–53.
- 53 Goldstein H, Models NM. With an application to discrete response data. *Biometrika* 1991;78:45–51.
- 54 Linkens AEMJH, Milosevic V, van der Kuy PHM, *et al.* Medication-Related hospital admissions and readmissions in older patients: an overview of literature. *Int J Clin Pharm* 2020;42:1243–51.
- 55 Berry JG, Gay JC, Joynt Maddox K, *et al.* Age trends in 30 day hospital readmissions: US national retrospective analysis. *BMJ* 2018;360:k497.



- 56 Kongkaew C, Hann M, Mandal J, *et al.* Risk factors for hospital admissions associated with adverse drug events. *Pharmacotherapy* 2013;33:827–37.
- 57 Thunander Sundbom L, Bingefors K. Women and men report different behaviours in, and reasons for medication non-adherence: a nationwide Swedish survey. *Pharm Pract* 2012;10:207–21.
- 58 Manteuffel M, Williams S, Chen W, *et al.* Influence of patient sex and gender on medication use, adherence, and prescribing alignment with guidelines. *J Womens Health* 2014;23:112–9.
- 59 Thunander Sundbom L, Hedborg K. Association between prescribed antidepressants and other prescribed drugs differ by gender: a nationwide register-based study in Sweden. *Nord J Psychiatry* 2019;73:73–9.
- 60 Silverstein MD, Qin H, Mercer SQ, *et al.* Risk factors for 30-day Hospital readmission in patients ≥65 years of age. *Proc* 2008;21:363–72.
- 61 Schwab C, Hindlet P, Sabatier B, *et al.* Risk scores identifying elderly inpatients at risk of 30-day unplanned readmission and accident and emergency department visit: a systematic review. *BMJ Open* 2019;9:e028302.
- 62 Jain S, Khara R, Mortensen EM, *et al.* Readmissions of adults within three age groups following hospitalization for pneumonia: analysis from the nationwide readmissions database. *PLoS One* 2018;13:e0203375.
- 63 Hasegawa K, Gibo K, Tsugawa Y, *et al.* Age-Related differences in the rate, timing, and diagnosis of 30-day readmissions in hospitalized adults with asthma exacerbation. *Chest* 2016;149:1021–9.
- 64 Shebeshi DS, Dolja-Gore X, Byles J. Unplanned readmission within 28 days of hospital discharge in a longitudinal population-based cohort of older Australian women. *Int J Environ Res Public Health* 2020;17:3136.
- 65 Sud M, Yu B, Wijeyesundera HC, *et al.* Associations between short or long length of stay and 30-day readmission and mortality in hospitalized patients with heart failure. *JACC Heart Fail* 2017;5:578–88.
- 66 Donzé J, Lipsitz S, Bates DW, *et al.* Causes and patterns of readmissions in patients with common comorbidities: retrospective cohort study. *BMJ* 2013;347:f1711.
- 67 Hijazi HH, Alyahya MS, Hammouri HM, *et al.* Risk assessment of comorbidities on 30-day avoidable Hospital readmissions among internal medicine patients. *J Eval Clin Pract* 2017;23:391–401.
- 68 Falvey JR, Bade MJ, Hogan C, *et al.* Preoperative activities of daily living dependency is associated with higher 30-day readmission risk for older adults after total joint arthroplasty. *Clin Orthop Relat Res* 2020;478:231–7.
- 69 Greysen SR, Stijacic Cenzer I, Auerbach AD, *et al.* Functional impairment and hospital readmission in Medicare seniors. *JAMA Intern Med* 2015;175:559–65.
- 70 Pickens S, Naik AD, Catic A, *et al.* Dementia and hospital readmission rates: a systematic review. *Dement Geriatr Cogn Dis Extra* 2017;7:346–53.
- 71 Chiang LY, Liu J, Flood KL, *et al.* Geriatric assessment as predictors of hospital readmission in older adults with cancer. *J Geriatr Oncol* 2015;6:254–61.
- 72 Butcher L. Oncologists seek to understand, address Hospital readmissions. *Oncology Times* 2016;38:1–10
- 73 Burhenn P, Sun C-L, Scher KS, *et al.* Predictors of hospital readmission among older adults with cancer. *J Geriatr Oncol* 2020;11:1108–14.
- 74 Leendertse AJ, Egberts ACG, Stoker LJ, *et al.* Frequency of and risk factors for preventable medication-related hospital admissions in the Netherlands. *Arch Intern Med* 2008;168:1890–6.
- 75 Leendertse AJ, Van Den Bernt PMLA, Poolman JB, *et al.* Preventable hospital admissions related to medication (HARM): cost analysis of the HARM study. *Value Health* 2011;14:34–40.
- 76 Garfinkel D, Bilek A. Inappropriate medication use and polypharmacy in older people. *BMJ* 2023;2020.
- 77 Tesfaye WH, Peterson GM, Castelino RL, *et al.* Medication regimen complexity and hospital readmission in older adults with chronic kidney disease. *Ann Pharmacother* 2019;53:28–34.
- 78 Fialová D, Onder G. Medication errors in elderly people: contributing factors and future perspectives. *Br J Clin Pharmacol* 2009;67:641–5.
- 79 Lavan AH, Gallagher PF, O'Mahony D. Methods to reduce prescribing errors in elderly patients with multimorbidity. *Clin Interv Aging* 2016;11:857–66.
- 80 Cook JA, Burke-Miller JK, Jonikas JA, *et al.* Factors associated with 30-day readmissions following medical hospitalizations among Medicaid beneficiaries with schizophrenia, bipolar disorder, and major depressive disorder. *Psychiatry Res* 2020;291:113168.